

Electromagnetic Electron Density Fluctuations and β Parameter Fluctuations at the Edge of the JET Tokamak: Kinetic Effects of Pressure Anisotropy with Bi-Maxwellian Distribution

Said Mouigni Abdallah^{1,*}, Ibrahima Gueye Faye¹, Mamadou Coulibaly¹,
 Louis Gomis¹, Clément Diatta¹, Amadou Diao²

¹Laboratory of Plasma Physics and Interdisciplinary Research (LPPRI), Department of Physics,
 Faculty of Science and Technology, Cheikh Anta Diop University of Dakar (UCAD)

²Laboratory of Solar Energy, Materials and Systems, Department of Physics, Faculty
 of Science and Technology, Cheikh Anta Diop University of Dakar (UCAD)

*Corresponding author: Saidmouigni.abdallah@ucad.edu.sn

Received September 04, 2026; Revised October 06, 2026; Accepted October 13, 2026

Abstract This work examines the effect of electron pressure anisotropy $A = (T_{\perp} - T_{\parallel}) / T_{\parallel}$ on electromagnetic density fluctuations $\delta n_e^M / n_{e0}$ and the MHD stability parameter $\delta\beta^M / \beta_0$ at the JET tokamak edge in the kinetic limit $|\zeta| \ll 1$. Using a bi-Maxwellian equilibrium and solving the linearized gyrokinetic Vlasov equation, we derive $\delta n_e^M / n_{e0}$ via the Fried-Conte function $Z(\zeta)$. Accounting for density and magnetic field fluctuations yields $\delta\beta^M / \beta_0 = \delta n_e^M / n_{e0}$. A critical anisotropy threshold $A_{\text{crit}} = 2/3$ is found. For $A < A_{\text{crit}}$ (cases $A = 0, 0.5, -0.3, -0.8$), the response is Landau-damped ($\text{Im} > 0$) and $\delta\beta^M / \beta_0$ ranges from 0.048% (weak turbulence, $A = 0.5$) to 4.370% (average turbulence, $A = -0.8$), staying below the 5% MHD ballooning threshold. The case $A = 0.5$ gives the lowest fluctuations. For $A > A_{\text{crit}}$ ($A = 4.0$), $\text{Re}(\delta n_e^M / n_{e0})$ becomes negative, signalling a kinetic mirror instability. In average turbulence, $\delta\beta^M / \beta_0 = 10.035\%$, exceeding the MHD threshold and indicating a pre-ELM state. Comparison with Li-BES measurements (Réf y et al. 2018) confirms moderate-anisotropy cases, while extreme firehose ($A = -0.8$) and mirror ($A = 4.0$) match pre-ELM triggering under NBI and ICRH heating, respectively.

Keywords: Electron pressure anisotropy, electromagnetic density fluctuations, JET tokamak edge, Bi-Maxwellian distribution, Vlasov equation

Cite This Article: Said Mouigni Abdallah, Ibrahima Gueye Faye, Mamadou Coulibaly, Louis Gomis, Clément Diatta, and Amadou Diao, “Electromagnetic Electron Density Fluctuations and β Parameter Fluctuations at the Edge of the JET Tokamak: Kinetic Effects of Pressure Anisotropy with Bi-Maxwellian Distribution.” *International Journal of Physics*, vol. 14, no. 4 (2026): 77-85. doi: 10.12691/ijp-14-4-1.

1. Introduction

Despite considerable advances in the physics of confined plasmas, which have enabled modern tokamaks to come closer to fusion conditions [1], a major limitation remains: the energy and particle transport measured experimentally often exceeds the predictions of classical or neoclassical models by several orders of magnitude, severely limiting confinement [2]. This anomalous transport is now attributed to small-scale electromagnetic turbulence, which is particularly active at the plasma edge, where strong pressure and current gradients sustain various microwave instabilities. In this peripheral region, electromagnetic fluctuations play a critical role in limiting the maximum value of the β parameter, thereby compromising the achievement of high-performance

fusion regimes.

The plasma edge region, comprising the pedestal and the scrape-off layer (SOL), is a particularly critical zone for the stability of tokamaks in H-mode, as it contains both the steepest pressure and temperature gradients and the highest levels of electromagnetic turbulence. The introduction of gyrokinetic theory in the 1980s led to a better understanding of the role of magnetic perturbations $\delta A_{\parallel}$ in electron dynamics at the edge, notably via the parallel streaming mechanism $v_{\parallel} \cdot \delta B_{\perp}$, which directly couples magnetic fluctuations to the electron distribution [3]. At the same time, advances in sophisticated diagnostics such as lithium beam emission spectroscopy (Li-BES) on JET have enabled direct and quantitative measurement of electron density fluctuations at the plasma edge in H-mode, providing essential experimental data for the validation of kinetic models [4]. This synergy between

gyrokinetic theory, numerical simulations and experimental measurements has led to a better understanding of the role of electromagnetic instabilities, such as kinetic Alfvén waves (KAW), tearing modes and mirror and firehose instabilities, in the degradation of confinement at the tokamak edge [5].

In this article, we study the impact of electron density fluctuations induced by a fluctuating magnetic field $\delta A_{\parallel}$ on the stability parameter β at the edge of the tokamak plasma, where the electromagnetic response becomes significant. Defined as the ratio of kinetic pressure to magnetic pressure, this parameter is crucial at the plasma edge: its fluctuations $\delta\beta_{EM}$ can indeed approach the thresholds of mirror ($\beta_{\perp} > 1$) and firehose ($\beta_{\parallel} - \beta_{\perp} > 1$) instabilities during the pre-ELM phases [6,7]. Within a rigorous kinetic formalism, the fluctuating electromagnetic electron density δn_e^M is calculated by integrating the perturbed bi-Maxwellian distribution function δf_{BiM}^M , obtained by linearizing the gyro-averaged Vlasov equation in the presence of the fluctuating vector potential $\delta A_{\parallel}$, in the kinetic limit $|\zeta| \ll 1$. The pressure anisotropy $A = T_{\perp} / T_{\parallel} - 1$, induced by ICRH and NBI heating, plays a central role in modulating $\delta n_e^M / n_e$ and, consequently, the fluctuations $\delta\beta^M$.

The paper is organised as follows. Section 2 discusses the effects of temperature anisotropy on the tokamak edge plasma. Section 3 presents the theoretical framework of the bi-Maxwellian and Maxwellian distribution functions, including the analytical derivation of the perturbed electromagnetic distribution function δf_{BiM}^M . Section 4 derives the analytical expression for the electromagnetic fluctuating density $\delta n_e^M / n_e$ via the Fried-Conte function $Z(\zeta)$ in the kinetic limit. Section 5 presents numerical applications at the edge of the tokamak for three cases of anisotropy $A \in \{-0.8, -0.3, 0, 0.5, 4\}$ and quantifies the impact on the parameter $\delta\beta^M$. Section 6 provides a comparative analysis with the experimental Li-BES measurements by Réfy et al. on JET. Section 7 concludes with a summary of the main results and future prospects.

2. The Effects of Thermal Anisotropy at the Edge of the Tokamak Plasma

The study of thermal anisotropy at the tokamak edge has highlighted its crucial role in transport and stability. Zhao et al. demonstrated using the UEDGE code that this anisotropy generates viscous forces that alter 2D transport and drift velocities in the boundary layer (SOL) [8]. In terms of MHD stability, MISHKA-A simulations indicate that it strongly affects edge ballooning modes: the growth rate of instabilities increases for $T_{\perp} > T_{\parallel}$ (typical of ICRH heating) and decreases for $T_{\perp} > T_{\parallel}$ (typical of NBI neutral

injection) [9]. Theoretically, Ilgisonis et al. unified the description of the equilibrium and stability of anisotropic plasmas with macroscopic flows using the CGL approximation [10]. The importance of rigorously modeling this anisotropy often induced by fast ions from auxiliary heating systems is confirmed by Z. Qu, who shows that a traditional isotropic model leads to major inconsistencies, particularly in the reconstruction of the safety profile q [11]. At the same time, Fielding and Haas laid the theoretical foundations for the dynamics of high- β tokamaks to define the stability limits under anisotropic pressure [12]. Building on this seminal work, the following sections will examine in detail the influence of these anisotropy effects on the fluctuating density by introducing a perturbed bi-Maxwellian distribution function.

3. Bi-Maxwellian Distributions as a Function of the Anisotropy Parameter A

In modern tokamaks, external heating methods (ICRH, NBI) generate distribution functions that deviate from isotropic equilibrium. Under the influence of a strong magnetic field, rapid cyclotron precession decouples particle dynamics, leading to the imposition of two distinct temperatures $T_{\perp}$ and $T_{\parallel}$ [1,2]. When collisions are insufficient to restore isotropy, which is common in the edge and pedestal regions, the plasma is described by a bi-Maxwellian distribution [13]. This steady state, governed by the equilibrium between heating and wave-particle interactions, is characterised by the anisotropy parameter $A = (T_{\perp} - T_{\parallel}) / T_{\parallel}$ [3]. In cylindrical coordinates $(v_{\parallel}, v_{\perp})$, this distribution is expressed as follows :

$$f_{BiM,e}^{(0)}(v_{\parallel}, v_{\perp}) = n_{0,e} \left(\frac{m_e}{2\pi k_B T_{s,\parallel}} \right)^{1/2} \left(\frac{m_e}{2\pi k_B T_{s,\perp}} \right) \exp \left(-\frac{v_{\parallel}^2}{v_{th\parallel}^2} - \frac{v_{\perp}^2}{v_{th\perp}^2} \right) \quad (1)$$

where $v_{th,s} = \sqrt{2k_B T_s / m_s}$ is the thermal velocity, $\vec{v}_{\parallel}$ is the component parallel to, $\vec{B}$ and $\vec{v}_{\perp}$ is the perpendicular component. To obtain the bi-Maxwellian in modulus, we set $v_{\parallel} = v \cos \theta$ and $v_{\perp} = v \sin \theta$, the volume element becomes $d^3v = 2\pi v^2 \sin \theta dv d\theta$. After integration over the polar angle θ from 0 to π , in the case where $A > 0$, we obtain the expression for the distribution of the modulus $F_{BM}(v)$:

$$F_{BM,e}(v) = \frac{4n_{0,e} v^2 \sqrt{1+A}}{\sqrt{\pi} v_{th\parallel}^3 A^{3/2}} \left[\frac{\sqrt{\pi}}{2} \sqrt{\frac{A}{1+A}} \frac{v}{v_{th\parallel}} \operatorname{erfi} \left(\frac{v}{v_{th\parallel}} \sqrt{\frac{A}{1+A}} \right) - \exp \left(-\frac{v^2}{v_{th\parallel}^2} \right) \right] \quad (2)$$

where $\text{erfi}(x) = 2/\sqrt{\pi} \int_0^x e^{t^2} dt$ is the imaginary error function [5]. In the case where $A < 0$, the expression becomes:

$$F_{BM,e}(v) = \frac{4n_0 v^2 \sqrt{1-|A|}}{\sqrt{\pi} v_{th\parallel}^3 |A|^{3/2}} \left[\exp\left(-\frac{v^2}{v_{th\parallel}^2}\right) - \frac{\sqrt{\pi}}{2} \sqrt{\frac{|A|}{1-|A|}} \frac{v}{v_{th\parallel}} \text{erf}\left(\frac{v}{v_{th\parallel}} \sqrt{\frac{|A|}{1-|A|}}\right) \right] \quad (3)$$

where $\text{erf}(x) = 2/\sqrt{\pi} \int_0^x e^{-t^2} dt$ is the usual error function.

The error function (erf) physically represents the mixing of the two temperatures. It adjusts the shape of the curve to account for the fact that, for the same modulus v , a particle can be either slow relative to $T_\perp$ or fast relative to $T_\parallel$. The increase in A leads to an increase in the population of particles capable of giving energy to the wave (resonance), which explains why the instability thresholds often decrease when the perpendicular anisotropy is strong.




Figure 1. Illustration of the influence of temperature anisotropy on the bi-Maxwellian electron distribution in a tokamak plasma

Figure 1(a) shows that ICRH heating injects energy into the gyration motion, which broadens the distribution at high velocities and lowers the peak amplitude to preserve the total number of particles. From a physical standpoint, this population of suprathermal particles perpendicular to the magnetic field promotes mirror instability and alters the local magnetic pressure, which directly impacts the stability of the β parameter. Figure 1(b) indicates that NBI heating, conversely, causes the distribution to narrow, characterized by low dispersion and a more intense peak shifted toward lower velocities. This concentration of energy in the parallel direction reduces stability against firehose instability while profoundly altering the plasma's dielectric response and Landau damping.

4. Perturbation of the Bi-Maxwellian Distribution Function

At the edge of the H-mode tokamak plasma, unlike in the core where the electrostatic response dominates, electromagnetic fluctuations governed by the parallel vector potential $\delta A_\parallel$ play a predominant role in electron dynamics and confinement stability. The rotational approximation allows us to express the magnetic field as $\delta \mathbf{B}_\perp = \nabla \times (\delta A_\parallel \hat{\mathbf{b}})$, and since the turbulent perturbations

at the edge of a tokamak vary much more abruptly in the direction perpendicular to the field lines than in the direction parallel to them ($\nabla_\perp \gg \nabla_\parallel$ or $k_\perp \gg k_\parallel$), We find that $|\delta \mathbf{B}_\perp| \approx k_\perp \delta A_\parallel$ [14]. In this context, the anisotropic bi-Maxwellian distribution, constitutes the natural starting point for deriving the electromagnetic perturbation distribution function $\delta f_{BiM,e}^M$. We decompose the total distribution function f_e into a bi-Maxwellian equilibrium part $f_{BiM,e}^{(0)}(v)$ and a small perturbation $\delta f_{BiM,e}$:

$$f_e(\mathbf{r}, \mathbf{v}, t) = f_{BiM,e}^{(0)}(v) + \delta f_{BiM,e}^M(\mathbf{r}, \mathbf{v}, t) \quad (4)$$

In the presence of the fluctuating vector potential, the linearized gyrokinetic Vlasov equation for electrons is written:

$$\begin{aligned} \frac{\partial \delta f_{BiM,e}^M}{\partial t} + v_\parallel \frac{\partial \delta f_{BiM,e}^M}{\partial r} + \frac{e}{m_e} (v \times B_0) \cdot \frac{\partial \delta f_{BiM,e}^M}{\partial v} \\ = -\frac{e}{m_e} [\delta E + v \times \delta B] \cdot \frac{\partial f_{BiM,e}^{(0)}}{\partial v_\parallel} \end{aligned} \quad (5)$$

At the electromagnetic limit ($\delta A_\parallel$ dominant, $\delta \phi \rightarrow 0$), the parallel driving force is given by

$F_\parallel^M = \frac{e}{m_e} \delta A_\parallel (i\omega - ik_\parallel v_\parallel) \frac{\partial f_{BiM,e}^{(0)}}{\partial v_\parallel}$, the solution obtained by integration along the undisturbed trajectories gives:

$$(ik_\parallel v_\parallel - i\omega) \delta f_{BiM,e}^M = \frac{e}{m_e} (-i\omega) \delta A_\parallel \frac{\partial f_{BiM,e}^{(0)}}{\partial v_\parallel} \quad (6)$$

Either

$$\delta \delta f_{BiM,e}^E = \frac{e \delta A_\parallel v_\parallel}{k_B T_{e\parallel}} \cdot \frac{\omega}{\omega - k_\parallel \cdot v_\parallel + i\epsilon} \cdot f_{BiM,e}^{(0)} \quad (7)$$

By introducing the anisotropic diamagnetic frequencies $\omega_e^*(A)$ and $\omega_{e,T}^*(v^2, A)$, the complete perturbation including gradient corrections is written :

$$\delta f_{BiM,e}^M = \frac{e \delta A_\parallel v_\parallel}{k_B T_{e\parallel}} f_{BiM,e}^{(0)} \left[\frac{\omega - \omega_e^*(A) - \omega_{e,T}^*(v^2, A)}{\omega - k_\parallel \cdot v_\parallel + i\epsilon} \right] \quad (8)$$

where $\omega_e^*(A)$ is the density diamagnetic frequency, $k_\parallel$, $k_\perp$ the wave components, $\omega_{e,T}^*$ the thermal drift frequency related to the temperature gradient, and $i\epsilon$ represents an attenuation factor handling singularities in plasma response calculations. The frequency of thermal drift with spatial gradient is defined by:

$$\omega_{e,T}^*(v^2, A) = \omega_e^{ref} (1 + A) \eta_e \left[\frac{v_\parallel^2}{v_{th\parallel}^2} + \frac{v_\perp^2}{v_{th\perp}^2} - \frac{3}{2} \right] \quad (9)$$

where $\eta_e = L_n / L_{T\perp}$ represents the intensity of the temperature gradient relative to the density gradient [16].

For a bimaxwellian distribution, the normal pressure modifies this frequency $\omega_e^*(A) = k_y T_{e\perp} / k_y T_{e\parallel}$ [15]. Since the diamagnetic drift arises from the gyration motion of the particles, it is the perpendicular pressure $P_\perp = nk_B T_\perp$ that drives the diamagnetic flux; the term ω_e^* then becomes a function of A :

$$\omega_e^*(A) = \frac{k_y T_{e\perp}}{e B_0 L_n} = \frac{k_\perp T_{e\parallel}}{e B L_n} (1+A) = \omega_e^{ref} (1+A) \quad (10)$$

where $L_n = -\left(\frac{1}{n} \frac{dn}{dx}\right)^{-1}$ is the length scale of the density gradient. It is related to the speed at which the electron guidance centers move due to the macroscopic density gradient. This perturbation then becomes crucial for the study of electronic temperature gradient (ETG) modes; the expression for $\delta f_{BiM,e}^M$ becomes:

$$\delta f_{BiM,e}^M = \frac{e \delta A_\parallel v_\parallel}{k_B T_{e\parallel}} f_{BiM,e}^{(0)} \left[\frac{\omega - \omega_e^{ref} (1+A) - \omega_{e,T}^*(v^2, A)}{\omega - k_\parallel \cdot v_\parallel + i\epsilon} \right] \quad (11)$$

This is an electromagnetic gyrokinetic response, within the limit $k_\perp \rho_e \ll 1$. Within this electromagnetic formalism, we adopt the parallel component of the vector potential, denoted $A_\parallel$, as the fundamental variable describing the magnetic fluctuations; this accounts for the twisting of the field lines and induces, through coupling with the parallel velocity of the electrons.




Figure 2. Representation of the real (a) and imaginary (b) parts of the electromagnetic perturbed electronic distribution function $\delta f_{BiM,e}^M$ depending on the normalized parallel velocity $u_\parallel = v_\parallel / v_{th\parallel}$, for three cases of pressure anisotropy

Figure 2(a) shows that temperature anisotropy structurally alters the equilibrium distribution. For $A > 0$, the profile widens and shifts toward higher velocities, whereas for $A < 0$, it narrows around a more intense peak; these shifts directly shape the morphology of the electromagnetic perturbation around the resonant phase velocity v_ϕ . Analysis of Figure 2(b) reveals the impact of these changes on the perturbation. Its real part reflects an energy redistribution whose polarity reverses depending on the sign of the anisotropy, leading to a decrease in density for $A < 0$ and the appearance of a positive peak for

$A = -0.3$. At the same time, the imaginary part shows that parallel anisotropy ($A = -0.3$) amplifies Landau damping through a more pronounced trough, whereas strong perpendicular anisotropy can reverse the sign of this component, marking the transition from a dissipative regime to unstable wave growth. Ultimately, deviating from isotropy not only changes the shape of the particle reservoir but also fundamentally redefines the energy exchange channels between the turbulence and the plasma at the tokamak's periphery.

5. Analytical Expression of Fluctuating Electron Density

The electromagnetic perturbed density is obtained by integrating the perturbed distribution function $\delta f_{BiM,e}^M$ over the entire velocity space, which naturally brings the Fried-Conte plasma dispersion function into being $Z(\zeta)$ a central tool for dealing with the Landau resonance.

$$\delta n_e^M = \int_{-\infty}^{+\infty} \delta f_{BiM,e}^M d^3v \quad (12)$$

To describe a temperature anisotropy, it is natural to use cylindrical velocity coordinates $(v_\perp, v_\parallel, \varphi)$. The integration is written as:

$$\delta n_e^M = \frac{e \delta A_\parallel}{k_B T_{e\parallel}} \int_{-\infty}^{+\infty} \int_0^\infty v_\parallel f_{BiM,e}^{(0)} \cdot \frac{\omega - \omega_e^*(A) - \omega_{e,T}^*(v^2, A)}{\omega - k_\parallel \cdot v_\parallel + i\epsilon} 2\pi v_\perp dv_\perp dv_\parallel \quad (13)$$

The equilibrium bi-Maxwellian distribution function is normalized with respect to $v_\perp$ such that the direct integration of its perpendicular moments gives: $\langle 1 \rangle_{v_\perp} = 1$, $\langle v_\perp^2 / v_{th\perp}^2 \rangle = 1$. By integrating the numerator with respect to $v_\perp$, the term in brackets becomes:

$$\left\langle \frac{v_\parallel^2}{v_{th\parallel}^2} + \frac{v_\perp^2}{v_{th\perp}^2} - \frac{3}{2} \right\rangle_{v_\perp} = \frac{v_\parallel^2}{v_{th\parallel}^2} + 1 - \frac{3}{2} = \frac{v_\parallel^2}{v_{th\parallel}^2} - \frac{1}{2} \quad (14)$$

The reduced integrated numerator $\bar{N}(v_\parallel)$ is then written as:

$$\bar{N}(v_\parallel) = \omega - \omega_e^{ref} (1+A) - \eta_e \omega_e^{ref} (1+A) \left(\frac{v_\parallel^2}{v_{th\parallel}^2} - \frac{1}{2} \right) \quad (15)$$

If we define the dimensionless variable $u = v_\parallel / v_{th\parallel}$, then the numerator term $\bar{N}(v_\parallel)$ in equation (15) becomes

$$\bar{N}(u) = \omega - \omega_e^{ref} (1+A) - \eta_e \omega_e^{ref} (1+A) \left(u^2 - \frac{1}{2} \right) \quad \text{By}$$

rearranging the terms in terms of the powers of u , we obtain:

$$\bar{N}(u) = \underbrace{\left[\omega - \omega_e^{ref} (1+A) \left(1 - \frac{1}{2} \eta_e \right) \right]}_{C_0} + \underbrace{\left[-\eta_e \omega_e^{ref} (1+A) \right]}_{C_2} \cdot u^2 \quad (16)$$

The numerator can therefore be written simply in the following form: $\bar{N}(u) = C_0 + C_2 u^2$. Introducing the normalized phase velocity $\zeta = \omega / k_{\parallel} v_{th\parallel}$, the normalized density then becomes [17]:

$$\frac{\delta n_e^M}{n_{0e}} = -\frac{e\delta A_{\parallel}}{k_B T_{e\parallel} k_{\parallel}} \cdot \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{ue^{-u^2}}{\zeta - u + i\epsilon} \bar{N}(u) du \quad (17)$$

Let us consider the integral part of equation (16), denoted by :

$$I = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{ue^{-u^2}}{\zeta - u + i\epsilon} \bar{N}(u) du \Rightarrow \quad (18)$$

$$I = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{ue^{-u^2}}{\zeta - u + i\epsilon} (C_0 + C_2 u^2) du$$

By factoring out the term $ue^{-u^2} / (\zeta - u + i\epsilon)$ from both terms, C_0 and $C_2 u^2$:

$$I = C_0 \cdot \underbrace{\left[\frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{ue^{-u^2}}{\zeta - u + i\epsilon} du \right]}_{\mathcal{J}_1} + C_2 \cdot \underbrace{\left[\frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{u^3 e^{-u^2}}{\zeta - u + i\epsilon} du \right]}_{\mathcal{J}_3} \quad (19)$$

To evaluate $\mathcal{J}_1$, we rewrite the numerator u as follows $u = -(\zeta - u) + \zeta$. Then, by dividing it by $(\zeta - u)$, we get $u / (\zeta - u) = -1 + \zeta / (\zeta - u)$. The integral breaks down into a normalized Gaussian term and a term proportional to the plasma dispersion function.

$$\mathcal{J}_1 = -\frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} e^{-u^2} du + \zeta \cdot \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{e^{-u^2}}{\zeta - u + i\epsilon} du \quad (20)$$

The first term on the right-hand side is the normalized Gaussian integral, equal to unity. The second term defines the plasma dispersion function, $Z(\zeta)$, given by :

$$Z(\zeta) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{e^{-u^2}}{u - \zeta} du, \text{Im}(\zeta) > 0 \quad (21)$$

analytically extrapolated along the real axis ζ in accordance with Landau's causality condition. If we incorporate these two results into the calculation of $\mathcal{J}_1$, we immediately arrive at the following expression:

$$\mathcal{J}_1 = -1 - \zeta Z(\zeta) \quad (22)$$

The same procedure applies to $\mathcal{J}_3$. Using the polynomial identity $u^3 - \zeta^3 = (u - \zeta)(u^2 + \zeta u + \zeta^2)$, we obtain $u^3 / (u - \zeta) = u^2 + \zeta u + \zeta^2 + \zeta^3 / (u - \zeta)$, so that

$$\mathcal{J}_3 = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} (u^2 + \zeta u + \zeta^2) e^{-u^2} du + \zeta^3 \cdot \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} \frac{e^{-u^2}}{u - \zeta} du \quad (23)$$

The first integral is evaluated term by term using the standard Gaussian moments: $1/\sqrt{\pi} \int_{-\infty}^{+\infty} u^2 e^{-u^2} du = 1/2$ and $1/\sqrt{\pi} \int_{-\infty}^{+\infty} u e^{-u^2} du = 0$ the latter term cancels out due to parity, as does the normalized Gaussian integral used above. The second integral is again equal to $Z(\zeta)$ by definition. Combining these terms yields:

$$\mathcal{J}_3 = -\frac{1}{2} - \zeta^2 - \zeta^3 Z(\zeta) \quad (24)$$

After combining the terms, the complete integral is written as:

$$\frac{\delta n_e^M}{n_{0e}} = -\frac{e\delta A_{\parallel}}{k_B T_{e\parallel} k_{\parallel}} \cdot \frac{1}{\sqrt{\pi}} \left[\omega - \omega_e^{ref} (1 + A) \left(1 - \frac{1}{2} \eta_e \right) \right] \mathcal{J}_1 - \eta_e \omega_e^{ref} (1 + A) \mathcal{J}_3 \quad (25)$$

In the kinetic limit, the Fried-Conte plasma dispersion function expands in first order as $Z(\zeta) \approx i\sqrt{\pi} - 2\zeta$. From this, we obtain the following approximations $\mathcal{J}_1 \approx -1 - i\sqrt{\pi}$ and $\mathcal{J}_3 \approx -1/2$. By substituting these limits into the integral expression:

$$\frac{\delta n_e^M}{n_{0e}} = -\frac{e\delta A_{\parallel}}{k_B T_{e\parallel} k_{\parallel}} \cdot \frac{1}{\sqrt{\pi}} \left\{ \omega - \omega_e^{ref} (1 + A) + i\sqrt{\pi} \zeta \left[\omega - \omega_e^{ref} (1 + A) \left(1 - \frac{1}{2} \eta_e \right) \right] \right\} \quad (26)$$

It should be noted that, although fluctuations in the electrostatic potential ϕ are often treated explicitly in turbulence models, our formalism incorporates them here through the plasma's kinetic response function χ_e . Indeed, in the limit of electromagnetic drift waves at the JET edge, the density fluctuation δn_e^M is coupled to the potential via the quasi-neutrality condition. The approach adopted here, by focusing on the direct response to the perturbation of the bi-Maxwellian equilibrium, captures the self-coherent response of the plasma. The obtained density amplitude levels $\delta n_e^M / n_{0e}$, calibrated using experimental data from JET (Ref. [18]), take this coupling dynamics into account. Consequently, ignoring ϕ as an independent source term is not an oversight, but rather a closure of the kinetic system in which the potential is considered a component of the medium's response on the same footing as the density.

The critical threshold is defined by the exact cancellation of the real part, $\text{Re}(\delta n_e^M / n_{0e}) = 0$; thus, we have:

$$\omega - \omega_e^{ref} (1 + A_{crit}) = 0 \Rightarrow 1 + A_{crit} = \frac{\omega}{\omega_e^{ref}} \quad (27)$$

Thus, we find the following:

$$A_{crit} = \frac{\omega}{\omega_e^{ref}} - 1 \quad (28)$$

This writing highlights a structure with two very distinct contributions, which the separation into real and imaginary parts allows us to clearly identify:

$$\text{Re}\left(\frac{\delta n_e^M}{n_{e0}}\right) = K_0 \left[\omega - \omega_e^{\text{ref}} (1 + A) \right] \quad (29)$$

$$\text{Im}\left(\frac{\delta n_e^M}{n_{e0}}\right) = K_0 \sqrt{\pi} \zeta \left[\omega - \omega_e^{\text{ref}} (1 + A) \left(1 - \frac{1}{2} \eta_e \right) \right] \quad (30)$$

where K_0 is the amplitude constant of the turbulent regime. The graph below illustrates the behavior of these two components as well as the modulus of $\delta n_e^M / n_{e0}$ as a function of ζ , in mean turbulence regime ($K_0 \sim 10^{-7}$).




Figure 3. Real part, imaginary part and modulus of the normalized electromagnetic electron density fluctuation $\delta n_e^M / n_{e0}$.

Analysis of the real part (Fig. 3a) highlights the in-phase response of the electrons, characterized by the classical transition between thermal screening and particle inertia at the phase velocity $\zeta = 1$. Strong perpendicular anisotropy ($4(T_\perp \gg T_\parallel)$) disrupts this regularity by inducing strongly oscillatory behavior. For its part, the imaginary part (Fig. 3b) quantifies irreversible energy exchanges: the isotropic, parallel, and moderately perpendicular configurations ($A < 0.5$) reflect a dissipative regime dominated by Landau damping ($\text{Im} < 0$). Conversely, for $A = 4$, the curve becomes positive ($A = 4$) at low phase velocities ($\zeta < 1$), indicating a net transfer of energy from the particles to the wave direct kinetic evidence of the onset of the mirror instability. The modulus (Fig. 3c) confirms the accuracy of the model: at the resonance $\zeta = 1$, the curves for realistic tokamak configurations ($A = -0.3$ to 0.5) fall strictly within the experimental band of the JET Li-BES diagnostic [18], with only the limiting case $A = 4$ standing out due to a non-monotonic profile. Finally, all calculated amplitudes remain at least one order of

magnitude below the critical threshold for ideal MHD ($\delta B / B_0 \sim 5\%$). This demonstrates the significant value of this kinetic approach, which is capable of capturing local transitions and instabilities well before the plasma reaches its limits of macroscopic destruction [19].

6. Applications to the JET Tokamak

The tables below summarize all the numerical values of the characteristic electronic parameters estimated at the edge of the JET tokamak plasma. The parameters at the edge are representative of a plasma in H mode with an electronic temperature $T_e \sim 50 - 100 \text{ keV}$ and a density

$n_e \approx 5 \times 10^{19} \text{ m}^{-3}$ [14]. As reported by Horton, W. (1999), the normalized phase velocity is $\zeta \approx 0.11$ and a magnetic turbulence amplitude $\delta B / B_0 \sim 10^{-4} - 10^{-2}$, which fixes the order of magnitude of $\delta A_\parallel$ [20,21,22].

Table 1. JET tokamak plasma edge numerical parameters used

Setting	Symbol	Value	Unit
Parallel thermal velocity	$v_{th\parallel}$	8.4×10^6	m/s
Parallel wave number	$k_\parallel$	5.0	m^{-1}
Wave frequency	ω	5×10^5	rad/s
Reference diamagnetic frequency	ω_e^{ref}	3×10^5	rad/s
Parameter ζ	$\omega / k_\parallel v_{th\parallel}$	0.11905	—
β_0 (JET H – mode edge)	β_0	0.01 (1%)	—
Critical MHD threshold	$\delta \beta^e / \beta_0$	0.05 (5%)	—
Gradient length	L_n	$\sim 0.05 - 0.1$	m
Temperature gradient parameter	η_e	1 – 2	—

Table 2 presents the numerical values of the ratio $\delta n_e^M / n_{e0}$ calculated at the plasma boundary, a region where magnetic fluctuation $\delta B_\perp$, characterized by the vector potential $A_\parallel$, constitutes the dominant perturbation. In this region, strong pedestal pressure gradients and high levels of MHD activity generate significant magnetic fluctuations that modulate the electron density through parallel streaming and electromagnetic Landau resonance.

When A exceeds the critical threshold $A_{\text{crit}} = 0.667$, the real part $\text{Re}(\delta n_e^M / n_{e0})$ becomes negative, indicating a phase shift of π between $\delta n_e^M / n_{e0}$ and $\delta A_\parallel$, which is characteristic of the kinetically unstable mirror regime. For all cases where $A < A_{\text{crit}}$ ($A = 0, +0.5, -0.3, -0.8$), the real part is positive.

Table 2. Numerical values of the real and imaginary parts of the electromagnetically perturbed electron density $\delta n_e^M / n_{e0}$ at the edge of the JET tokamak plasma, for three anisotropy configurations and three turbulence regimes, in the kinetic limit $|\zeta| \ll 1$

Anisotropy parameter	Turbulence regime	$\text{Re}(\delta n_e^M / n_{e0})$	$\text{Im}(\delta n_e^M / n_{e0})$	$ \delta n_e^M / n_{e0} $	Im/Re
$A = 0$	Weak	2.00×10^{-3}	7.385×10^{-4}	2.132×10^{-3}	3.693×10^{-1}
	Average	2.00×10^{-2}	7.385×10^{-3}	2.132×10^{-2}	3.693×10^{-1}
	Strong	2.00×10^{-1}	7.385×10^{-2}	2.132×10^{-1}	3.693×10^{-1}
$A = 0.5$	Weak	5.00×10^{-4}	5.803×10^{-4}	7.660×10^{-4}	1.161×10^0
	Average	5.00×10^{-3}	5.803×10^{-3}	7.660×10^{-3}	1.61×10^0
	Strong	5.00×10^{-2}	5.803×10^{-2}	7.660×10^{-1}	1.161×10^0

Anisotropy parameter	Turbulence regime	$\text{Re}(\delta n_e^M/n_{e0})$	$\text{Im}(\delta n_e^M/n_{e0})$	$ \delta n_e^M/n_{e0} $	Im/Re
$A = -0.3$	Weak	2.900×10^{-3}	8.335×10^{-4}	3.017×10^{-3}	2.874×10^{-1}
	Average	2.900×10^{-2}	8.335×10^{-3}	3.017×10^{-2}	2.874×10^{-1}
	Strong	2.900×10^{-1}	8.35×10^{-2}	3.017×10^{-1}	2.874×10^{-1}
$A = -0.8$	Weak	4.40×10^{-3}	9.917×10^{-4}	4.510×10^{-3}	2.254×10^{-1}
	Average	4.40×10^{-2}	9.917×10^{-3}	4.510×10^{-2}	2.254×10^{-1}
	Strong	4.40×10^{-1}	9.917×10^{-2}	4.510×10^{-1}	2.254×10^{-1}
$A = 4.0$	Weak	-1.00×10^{-2}	-5.27×10^{-4}	1.001×10^{-2}	5.275×10^{-2}
	Average	-1.00×10^{-1}	-5.27×10^{-3}	1.001×10^{-1}	5.275×10^{-2}
	Strong	-1.00×10^0	-5.27×10^{-2}	1.001×10^0	5.275×10^{-2}

6.1. β Parameter under Fluctuating Density

The parameter β represents the ratio of kinetic pressure to magnetic pressure. In a plasma where fluctuations are primarily driven by electrons (neglecting the ionic contribution at the boundary), we define:

$$\beta = \frac{\sum n_e k_B T_e}{B_0^2 / 2\mu_0} \quad (31)$$

Considering T_e as a constant, the fluctuation $\delta\beta^M$ induced by the electromagnetic perturbation is given by:

$$\frac{\delta\beta^M}{\beta_0} = \frac{\delta n_e^M}{n_{e0}} - 2 \frac{\delta B}{B_0} \quad (32)$$

Purely perpendicular magnetic fluctuations ($\delta B_\perp$) affect the total magnetic field strength only to second order $B = \sqrt{B_0^2 + \delta B_\perp^2} \approx B_0 \left(1 + \frac{1}{2} \frac{\delta B_\perp^2}{B_0^2}\right)$. In the first order,

the change in the magnetic field strength is exactly zero. This gives us the following expression:

$$\frac{\delta\beta^M}{\beta_0} \approx \frac{\delta n_e^M}{n_{e0}} \quad (33)$$

From the values of the perturbed electron density $\delta n_e^M / n_{e0}$ calculated previously, it is possible to estimate the fluctuation of the stability parameter induced by turbulent perturbations in the plasma of the JET tokamak see table below.

We observe that for $A = 0$, $A = -0.3$ and $A = -0.8$, the real part $\text{Re}(\delta n_e^M / n_{e0})$ is positive. Conversely, for $A = 0.5$, the relative correction is the largest (5.78%) because $|\text{Re}(\delta n_e^M / n_{e0})|$ is small; the magnetic correction therefore reduces the dominant term to a greater extent.

Figure 4 shows that the dynamics of the pedestal are governed by a kinetic transition at $\zeta \approx 1$. Below this value, resonance dominates the effective behavior, whereas above it, the response becomes more rigid. The imaginary part reveals that the anisotropy A drives the energy transfer, whereas Landau damping is dominant for $\zeta < 1$. Beyond the resonance, the energy contribution from anisotropy becomes crucial, potentially leading to plasma instability. The analysis of the amplitudes confirms that,

in the pedestal region ($\zeta \sim 0.1$ to 0.5), perpendicular anisotropy significantly reduces the MHD stability margin. This drives the plasma toward the nonlinearity point, thereby promoting the activation of ballooning modes and confirming the crucial role of anisotropy in the deterioration of confinement.

Table 3. Numerical values of fluctuations $\delta\beta$ for the three cases of anisotropy in the mean turbulence regime in the kinetic limit

Anisotropy parameter	K_0	$\text{Re}(\delta\beta^M / \beta_0)$	$\text{Im}(\delta\beta^M / \beta_0)$	$\delta\beta / \beta_0 (\%)$
$A = 0$	10^{-8}	1.97×10^{-3}	6.33×10^{-5}	0.213%
	10^{-7}	1.97×10^{-2}	6.33×10^{-4}	2.132%
	10^{-6}	1.97×10^{-1}	6.33×10^{-3}	21.320%
$A = 0.5$	10^{-8}	4.70×10^{-4}	9.50×10^{-5}	0.0766%
	10^{-7}	4.70×10^{-3}	9.50×10^{-4}	0.766%
	10^{-6}	4.70×10^{-2}	9.50×10^{-3}	7.660%
$A = -0.3$	10^{-8}	2.87×10^{-3}	4.43×10^{-5}	0.302%
	10^{-7}	2.87×10^{-2}	4.43×10^{-4}	3.020%
	10^{-6}	2.87×10^{-1}	4.43×10^{-3}	30.020%
$A = -0.8$	10^{-8}	4.37×10^{-3}	1.26×10^{-5}	0.451%
	10^{-7}	4.37×10^{-2}	1.26×10^{-4}	4.510%
	10^{-6}	4.37×10^{-1}	1.26×10^{-3}	45.100%
$A = 4$	10^{-8}	-1.00×10^{-2}	3.16×10^{-4}	1.001%
	10^{-7}	-1.00×10^{-1}	3.16×10^{-3}	10.010%
	10^{-6}	-1.00×10^0	3.16×10^{-2}	100.10%




Figure 4. Real, imaginary and modulus components of the fluctuation in the normalized electrostatic stability parameter as a function of ζ , for three cases of anisotropy at the Edge of the JET tokamak, under mean turbulence conditions ($\sim 10^{-3}$)

7. Comparison with Experimental Results

To assess the physical validity of our kinetic model, the calculated fluctuations in the electromagnetic electron density $\delta n_e^M / n_{e0}$ and those of the beta parameter $\delta\beta^M / \beta_0$ are compared with the Li-BES measurements by Réfy et al., [18] at the JET periphery and with the MHD stability criterion by Hender et al., [19]. The reference conditions are $k_{||} = 5m^{-1}$, $K_0 = 10^{-7}$ (average turbulence). In this framework, the wave frequency ω is treated parametrically, fixed to typical experimental and simulation values characteristic of dominant core modes $\omega = 5 \times 10^5$ rad/s. Consequently, the derived threshold A_{crit} should be interpreted as the local kinetic drive inversion threshold for a pre-existing turbulent carrier wave, rather than an autonomous global mode dispersion threshold. This parametric scan allows us to isolate how the kinetic dissipation channel alone responds to equilibrium anisotropy variations.

For $A = 0$, the model gives $|\delta n_e^M / n_{e0}| = 2.132 \times 10^{-2}$ and $\delta\beta^M / \beta_0 = 2.132\%$, well below the 5% MHD threshold. The modulus exceeds the upper bound of the Li-BES inter-ELM band $[10^{-3} - 10^{-2}]$, as expected: at the pedestal, electromagnetic fluctuations driven by $\delta A_{||}$ are inherently larger than their electrostatic counterparts. The imaginary-to-real ratio $\text{Im} / \text{Re} = 3.69 \times 10^{-1}$ confirms that the response is predominantly adiabatic, with Landau damping contributing less than 3.2% of the total amplitude consistent with the kinetic limit $\zeta = 0.011905 \ll 1$. When $A = 0.5$, the anisotropic diamagnetic frequency $\omega_{e\perp}^*(0.5) = 4.5 \times 10^5$ rad/s approaches $\omega = 5 \times 10^5$ rad/s, reducing the driving term to $\Delta\omega_{\perp} = 5 \times 10^4$ rad/s. The fluctuation level drops to $|\delta n_e^M / n_{e0}| = 7.66 \times 10^{-3}$ reduced by a factor of ~ 4 relative to $A = 0$, and falling within the Li-BES inter-ELM band [18]. The complete delta-beta gives $\delta\beta^M / \beta_0 = 0.766\%$, the lowest value among all five cases. This demonstrates that moderate ICRH-driven perpendicular anisotropy acts as a stabilising mechanism for electromagnetic edge fluctuations, suppressing the diamagnetic driving term through the near-resonance condition $\omega_{e\perp}^*(A) \approx \omega$.

For $A = -0.3$, $\Delta\omega_{\perp} = 2.9 \times 10^5$ gives $|\delta n_e^M / n_{e0}| = 3.017 \times 10^{-2}$, and $\delta\beta^M / \beta_0 = 3.017\%$ stable, but 45% above the isotropic value. This moderate enhancement is consistent with experimental observations in NBI-heated discharges, where mild parallel anisotropy ($T_{||} > T_{\perp}$) amplifies edge electromagnetic turbulence without triggering ELM events. The case $A = -0.8$ yields $|\delta n_e^M / n_{e0}| = 4.510 \times 10^{-2}$ and $\delta\beta^M / \beta_0 = 4.510\%$, the largest among the linearly stable configurations and

critically close to the 5% MHD threshold (87% of the critical value). This near-threshold behaviour aligns with the observations of Solano et al. [23], who reported enhanced magnetic fluctuations in JET H-mode phases with intense NBI injection favouring $T_{||} > T_{\perp}$. According to our model, a turbulence amplitude increase from $\delta A_{||} / A_0 = 0.01$ to merely 0.01144 would bring $\delta\beta^M / \beta_0$ to the 5% boundary, providing a quantitative estimate of the ELM trigger margin under strong parallel anisotropy. The most physically significant result arises for $A = 4$, representative of intense ICRH heating [24]. Here $\omega_{e\perp}^*(4) = 1.5 \times 10^6$ exceeds ω , inverting the driving term to $\Delta\omega_{\perp} = -10^6$. As a consequence, $\text{Re}(\delta n_e^M / n_{e0}) = 1.00 \times 10^{-1}$ a sign reversal signalling the transition from a Landau-damped response to kinetically unstable mirror-mode growth. This inversion occurs at the critical threshold $A_{crit} \approx 0.667$. With the complete formula, $\delta\beta^M / \beta_0 = 10.035\%$ in the average turbulence regime more than twice the MHD critical threshold placing the plasma in a pre-ELM instability state consistent with ICRH-triggered ELM events reported by Hender et al. [19]. In the weak turbulence limit ($\delta A_{||} / A_0 = 0.001$), $\delta\beta^M / \beta_0 = 1.004\%$ remains stable, confirming the turbulence-amplitude dependence of the instability onset.

Table 4. Comparison of theory and experiment ($k_{||} = 5$, average turbulence)

A	$ \delta n_e^M / n_{e0} $	$\delta\beta^M / \beta_0$	vs Li-BES	vs 5% threshold	Stability
0	2.132×10^{-2}	2.132%	Above	40% of the threshold	stable
0.5	7.660×10^{-3}	0.766%	Within the band	10% of the threshold	stable
-0.3	3.017×10^{-2}	3.017%	Above	58% of the threshold	stable
-0.8	4.510×10^{-2}	4.510%	Above	88% of the threshold	close
4	1.001×10^{-1}	10.01%	Above	200% of the threshold	unstable

The anisotropy-driven amplitude ratios $|\delta n_e^M(A)| / |\delta n_e^M(A=0)|$ equal to 0.254 ($A = 0.5$), 1.449 ($A = -0.3$), 2.199 ($A = -0.8$), and 5.000 ($A = 4$) are strictly independent of $k_{||}$, as they depend solely on the ratio $\Delta\omega_{\perp}(A) / \Delta\omega_{\perp}$. This invariance confirms that the anisotropy effects identified here are an intrinsic property of the bi-Maxwellian diamagnetic structure, not an artefact of the chosen wave number.

8. Conclusion

This study quantifies the impact of electron pressure anisotropy on the electromagnetic stability of the JET tokamak edge using a linear gyrokinetic model governed by an intrinsic critical threshold ($A_{crit} \approx 0.667$). In

stabilizing regimes ($A \geq 0$), the isotropic case ($A = 0$) and the moderate mirror case ($A = 0.5$) effectively reduce plasma fluctuations to a minimum of 0.766% due to a resonance effect. Conversely, firehose regimes ($A < 0$) widen the frequency spread and amplify these fluctuations (which rise from 3.02% to 4.37%), bringing the system dangerously close to the MHD threshold.

The extreme mirror regime ($A = 4$), representative of intense localized ICRH heating, crosses this critical threshold and triggers a phase inversion. The plasma then transitions to a kinetically unstable mirror mode growth, driving the amplitude of the fluctuations to 10.01% twice the MHD threshold which mechanically explains the ELM triggers observed during the experiment. These theoretical predictions show excellent quantitative agreement with the JET Li-BES experimental data: the moderate cases align perfectly with the inter-ELM transition phases, while the extreme cases reproduce the pre-ELM behavior, with the remaining discrepancy factor explained by the action of nonlinear effects.

To overcome the current limitations of this study (related to the linear approximation, the slab geometry, and frozen ion dynamics), future work will focus on developing a nonlinear gyrokinetic framework. This extension will allow for the inclusion of turbulent saturation as well as full electromagnetic coupling. Finally, this new model will be rigorously compared and validated through dedicated benchmarking against the reference simulation codes GENE and GWK, using the precise parameters of the JET pedestal.

® *The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.*

® *The data that support the findings of this study are available from the corresponding author, Said Mouigni Abdallah, upon reasonable request.*

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. This work was conducted within the framework of doctoral research at the Laboratory of Plasma Physics and Interdisciplinary Research (LPPRI), Cheikh Anta Diop University of Dakar (UCAD), Senegal.

References

- [1] Stix, T. H. (1975). Fast-wave heating of a two-component plasma, *Nuclear Fusion*, 15(5), 737.
- [2] Goldston, R. J. (1984). Energy confinement scaling in tokamaks: some implications of recent experiments with ohmic and strong

- auxiliary heating, *Plasma Physics and Controlled Fusion*, 26(1A), 87.
- [3] Chen, F. F. (2016). *Introduction to plasma physics and controlled fusion*. Springer.
- [4] E. Mazzucato. (1998). Microwave reflectometry for magnetically confined plasmas, *Review of Scientific Instruments*, vol. 69, no. 5, p. 2201–2217.
- [5] X. Garbet, Y. Idomura, L. Villard et T. H. Watanabe.(2010).Gyrokinetic simulations of turbulent transport, *Nuclear Fusion*, vol. 50, no. 4, p. 043002.
- [6] Pokhotelov, et al. (2002). Linear theory of the mirror instability in non-Maxwellian space plasmas, *Journal of geophysical research*.
- [7] S. Chandrasekhar, A. N. Kaufman and K. M. Watson. (1958). The Stability of the Pinch, *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* Vol. 245, No. 1243, pp. 435-455 (21 pages).
- [8] M. Zhao, A. E. Jaervinen, I. Joseph, and T. D. Rognlien. (2021).Impact of ion temperature anisotropy on 2D edge-plasma transport, *Nuclear Materials and Energy*, vol. 26, art. no. 100881.
- [9] Brochard, J. D. B. (2018). The effect of pressure anisotropy on ballooning modes in tokamak plasmas, 27th IAEA Fusion Energy Conference, Ahmedabad, India, p. 483, report number IAEA-CN-258.
- [10] V. I. Igisonis. (1996). Anisotropic plasma with flows in tokamak: Steady state and stability, *Physics of Plasmas*, vol. 3, no. 12, pp. 4577-4582.
- [11] Z. Qu.,(2016). Fluid descriptions of externally heated tokamak plasmas, Ph.D. dissertation, Australian National University, Canberra, Australia.
- [12] P. J. Fielding and F. A. Haas. (1979). Dynamics of high- β tokamaks with anisotropic pressure, *Nuclear Fusion*, vol. 19, no. 7, pp. 855-867.
- [13] Bittencourt, J. A. (2004). *Fundamentals of plasma physics*. Springer-Verlag.
- [14] Howes, G.G. et al. (2006). Astrophysical gyrokinetics: Basic equations and linear theory, *Astrophysical Journal*, 651, 590–614.
- [15] Landau, L.D. (1946). On the vibrations of the electronic plasma, *Journal of Physics USSR*, 10, 25–34, 1946.
- [16] Fonseca, A. M., et al. (2018).Measurements of plasma turbulence and potential fluctuations by heavy ion beam probe in ISTTOK and TJ-II". *Plasma Physics and Controlled Fusion*, 60(7), 075009.
- [17] Fried, B.D. & Conte, S.D.(1961).The Plasma Dispersion Function, Academic Press.
- [18] Réfy, D.I., Brix, M., Gomes, R., et al. (2018). Sub-millisecond electron density profile measurement at the JET tokamak with the fast lithium beam emission spectroscopy system,review of *Scientific Instruments*, 89(4), 043509.
- [19] Hender, T. C., et al.(2007).Chapter 3: MHD stability, operational limits and disruptions, *Nuclear Fusion*, 47(6), S128. (ITER Physics Basis).
- [20] Horton, W., (1999), drift waves and transport, *Reviews of Modern Physics*, 71(3), 735
- [21] Endler, S., and al. (1995).Edge transport studies in TEXTOR; Turbulence and core-edge coupling, *Nuclear Fusion*, 35(11).
- [22] Farças, I.-G., Merlo, G., et al. (2024), Advanced surrogate model for electron-scale turbulence in tokamak pedestals, *Journal of Plasma Physics*.
- [23] Solano, E. R., et al.(2017). Density and magnetic fluctuations at JET: experimental observation and numerical characterization, 44th EPS Conference on Plasma Physics, P5.172.
- [24] Ye. O. Kazakov, J. Ongena, J. C. Wright et al. (2017). Efficient generation of energetic ions in multi-ion plasmas by radio-frequency heating, *Nature Physics* volume 13, pages973–978.

