

Combined Application of Biological-Photocatalytic Process in Degradation of Reactive Black Dye: An Excellent Outcome

Maulin P Shah*

Industrial Waste Water Research Laboratory, Applied & Environmental Microbiology Lab, Enviro Technology Limited (CETP), Ankleshwar, Gujarat, India

*Corresponding author: shahmp@uniphos.com

Received September 24, 2013; Revised November 23, 2013; Accepted December 05, 2013

Abstract In the present study, the decolorization and degradation of Reactive Black 5 azo dye was investigated by biological, photocatalytic (UV/TiO₂) and combined processes. An application of *Pseudomonas aeruginosa* ETL-2211 in treatment of the synthetic medium containing RB5 indicated complete decolorization of the dye with 200 mg/L in less than 24 h. Degradation of the aromatic rings, resulting from the destruction of the dye, did not occur during the biological treatment. Mineralization of 50 mg/L RB5 solution was obtained after 80 min by photocatalytic process (in presence of 0.2 g/L TiO₂). COD (chemical oxygen demand) was not detectable after complete decolorization of 50 mg/L RB5 solution. However, photocatalytic process was not effective in the removal of the dye at high concentrations (≥ 200 mg/L). With 200 mg/L concentration, 74.9% of decolorization was achieved after 4 h illumination under photocatalytic process and the absorbance peak in UV region (attributed to aromatic rings) was not completely removed. A two-step treatment process, namely, biological treatment by bacteria followed by photocatalytic degradation, was also assessed. In the combined process (with 200 mg/L RB5), absorbance peak in UV region significantly disappeared after 2 h illumination and about 60% COD removal was achieved in the biological step. It is suggested that the combined process is more effective than the biological and photocatalytic treatments in the remediation of aromatic rings.

Keywords: azo dyes, *Pseudomonas aeruginosa*, combined wastewater treatment

Cite This Article: Maulin P Shah, "Combined Application of Biological-Photocatalytic Process in Degradation of Reactive Black Dye: An Excellent Outcome." *American Journal of Microbiological Research* 1, no. 4 (2013): 92-97. doi: 10.12691/ajmr-1-4-5.

1. Introduction

Azo dyes, representing one group of common textile dyes, contain one to four azo groups usually attached to two radicals of which at least one, but usually both, is aromatic groups. These molecules are chemically stable and difficult to biodegrade aerobically [1]. More than 100,000 commercial dyes are available to textile industries with over 700,000 tons of commercial dye being produced annually. The reactive azo dyes are highly recalcitrant to conventional wastewater treatment processes. In fact, as much as 90% of reactive dyes can remain unaffected after activated sludge treatment [2]. An estimated 10-15% of the synthetic dyes are discharged into wastewater as effluent which carries the potential to cause environmental damage [3]. In many cases, colored textile wastewater is aesthetically unacceptable, hinders light penetration, damages the quality of the receiving streams, and may be inhibitory towards biological wastewater treatment systems, to food chain organisms and to aquatic life. Consequently, it is important to remove dyestuff from the wastewater before discharging it to the environment [4].

The strong electron-withdrawing groups characterizing the chemical structure of dyes protect them from the bacterial oxygenases [5]. Therefore, conventional activated sludge systems have difficulty in handling such wastewater, particularly in removing the color [4]. The traditional treatment techniques applied in textile wastewater, such as chemical coagulation/flocculation, membrane separation (ultrafiltration, reverse osmosis) or activated carbon adsorption are known to be costly and only transfer the pollutants from the liquid to the solid phase [4,6]. Therefore, more efficient methods for treatment of dyed effluents should be examined. Most azo dye degrading microorganisms cleave the azo bond(s) of the respective azo dye and colorless, possibly toxic, aromatic amines are formed [7]. Thus, an alternative method must be considered for oxidization of the toxic aromatic amines. Advanced oxidation processes (AOPs) produce a highly reactive non-specific oxidant called hydroxyl radicals (OH•), capable of destroying a wide range of organic pollutants in water and wastewater [2,8]; presently, destruction of organic compounds using TiO₂/UV photocatalytic process is one of the various advanced oxidation processes applied. Numerous advanced treatment technologies have been developed to reduce

contaminants from textile wastewater. To effectively treat such wastewater, often a combined chemical and biological oxidation process is required as purely biological treatment is usually very slow or not possible at all [4]. A lot of information is available on removal and degradation of Remazol Black B using pure strains of bacteria, fungi, algae and yeast such as *Shewanella* strain J18 143, *Rhizopus arrhizus*, *Phlebia tremellosa*, *Trametes versicolor*, *Bjerkandera* sp. BOS55, *P. chrysosporium*, *Chlorella vulgaris*, *Candida tropicalis*, *Saccharomyces cerevisiae*, *Kluyveromyces marxianus* IMB3, *P. mirabilis*, *P. luteola*, *Pseudomonas* sp. and *K. rosea* [9,10,11,12]. The aim of the present study was to investigate the efficiency of photocatalytic process as a post-treatment stage in the removal of aromatic rings produced through biological treatment.

2. Materials and methods:

2.1. Chemicals and Dyes

Azo dye, Reactive Black 5 (RB5), was obtained from a local company and used without further purification. The dye is widely used by textile industries in Iran. Figure 1 shows the chemical structure of RB5. TiO₂-p25 was purchased from Germany. It has a specific surface area (BET) of approximately 50 m²/g and an average particle size of 21 nm.




Figure 1. Chemical structure of Reactive Black 5

2.2. Bacteria and Growth Medium

The bacterial strain used was isolated from dye wastewater obtained from final effluent treatment plant (FETP) of Ankleshwar, Gujarat, India. The principle of sequential selective enrichment batch culture for selection of dye decolorizing bacteria was employed in synthetic wastewater medium (SWM) with Basic Violet 3 as the carbon source. The basic composition of the synthetic wastewater medium was (g/L); (NH₄)₂SO₄ 0.28, NH₄Cl 0.23, KH₂PO₄ 0.067, MgSO₄ · 7H₂O 0.04, CaCl₂ · 2H₂O 0.022, FeCl₃ · 6H₂O 0.005, NaCl 0.15, NaHCO₃ 1.0, and 1 mL/L of a trace element solution containing (g/L); ZnSO₄ · 7H₂O 0.01, MnCl₂ · 4H₂O 0.1, CuSO₄ · 5H₂O 0.392, CoCl₂ · 6H₂O 0.248, NaB₄O₇ · 10H₂O 0.177, and NiCl₂ · 6H₂O 0.02. The textile wastewater used for isolation of dye-decolorizing bacteria was acclimatized for 8 weeks prior to transfer into 250mL Erlenmeyer flasks containing 100mL SWM. After incubation of the flasks, a mixed culture that showed quick and stable decolorization activity was transferred to newly prepared SWM. After

five successive transfers, it was plated on SWM agar containing 20 mg/L of each dye and incubated at 30°C for 5 days. Bacterial colonies around which clear zones expanded quickly were picked for further studies and designated as ETL 1-5. To check for the dye-degrading potential of each bacterial isolate, preliminary batch experiments were carried out using sterile 250mL Erlenmeyer flasks containing 100mL of SWM spiked with dye, after which the solution was inoculated with freshly grown bacterial cells. The final pH was 7.2. The bacterial isolate that showed the highest ability to degrade the triarylmethane dyes was selected and used for subsequent investigations. The selected bacterium (ETL-1) was characterized and identified as *Pseudomonas aeruginosa* using Gram stain, spore test, motility test, and a battery of biochemical and physiological tests as described by Vanderzant and Splittstoesser [13] and Cheesbrough [14] and with reference to the Bergey's Manual of Determinative Bacteriology [15].

2.3. Biological Experiments and Decolorization Procedures

Examination of decolorization and biodegradation ability of *Pseudomonas aeruginosa* was carried out in synthetic medium containing: (glucose, 10; Yeast extract, 0.34; NH₄Cl, 0.84; KH₂PO₄, 0.134; K₂HPO₄, 0.234; MgCl₂·6H₂O, 0.084) g/L and 200 mg/L RB5. The pH of the solution was adjusted to 7±0.2 before being autoclaved. A 5% (v/v) of the culture suspension in 0.85% saline was inoculated into 100 mL of medium in 250-mL flasks and incubated in shaking incubator at 150 rpm and at 32°C for 72 h. In addition, control flasks containing the same medium without inoculums were also kept to observe the abiotic decolorization. At regular time intervals, an aliquot (4 mL) of the culture media was withdrawn from the flasks, centrifuged at 12000 rpm for 10 min and the supernatant was analyzed for the remaining dye concentration. Decolorization of dye was determined by measuring the absorbance of culture supernatant at λ_{max} (597 nm). The percentage of decolorization was calculated according to the following formulation:

$$\text{Percentage of decolorization} = (A_i - A_t) / A_i \times 100 \quad (1)$$

Where A_i and A_t are the initial absorbance of the dye solution and at sampling time (t) after inoculation, respectively [16]. Spectral scanning between 200 and 800 nm using a UV-vis spectrophotometer (Shimadzu, Japan) was performed in order to analyze the dye degradation process. In this study, absorbance measurement at 254 nm wavelength was used to indicate the presence of aromatic compound in the sample [17].

2.4. Photocatalytic Experiments

Photocatalytic reactor and light source Photocatalytic experiments were performed in a batch system (pyrex vessel) with a total capacity of 1 L (batch mode). The total volume of the dye solution in the system was 800 mL. The reactor containing reaction solution (mixture of dye solution and photocatalyst powder) was placed on a magnetic stirrer to provide appropriate mixing. A medium-pressure mercury lamp (UV-C, maximum emission at 247.3 nm) with a power of 150 W and 66 mm

length (manufactured by ARDA, France) was used as the light source. The UV lamp was placed centrally and parallel to the length of the cylinder in the reactor, inside a quartz tube [18,19]. The reactor was covered with an aluminum foil sheet to prevent UV emission around it. Due to the production of heat by the light source, the temperature of the solution was maintained constant at $21 \pm 3^\circ\text{C}$ in all experiments by cooling water around the vessel.

2.5. Analytical Procedures

In order to perform the photocatalytic experiments, the working solution, containing 50 mg/L of dye and 0.2 g/L of TiO_2 was prepared using tap water. Prior to the photocatalytic process, the suspensions were magnetically stirred in the dark for 30 min to reach adsorption-desorption equilibrium between the dye and TiO_2 . Samples (5 mL) were taken at specific time intervals, centrifuged at 6000 rpm for 15 min and then filtered through a $0.45 \mu\text{m}$ syringe filter in order to separate TiO_2 particles [20]. Measurement of absorbance was carried out at maximum wavelength of RB5 ($\lambda_{\text{max}} = 597 \text{ nm}$). UV-vis spectra of samples were recorded between 200 and 800 nm. Photocatalytic degradation of different concentrations of RB5, namely, 100, 200, 300 and 500 mg/L was also examined.

2.6. Combined Biological-photochemical Process

To carry out these experiments, synthetic medium was treated with *Pseudomonas aeruginosa* ETL-2211 cells for 24 h and after centrifugation and filtration, fed into the reactor. At different time intervals, samples were withdrawn, centrifuged and filtered through a $0.45 \mu\text{m}$ membrane filter following which UV-vis scan spectroscopy and COD measurements were carried out. COD measurements (for both biological and photocatalytic steps of the process) were performed following the closed reflux, colorimetric method using a COD reactor (Hack, USA), according to standard procedures [21].

3. Results

Biological experiments *Pseudomonas aeruginosa* ETL-2211 was able to remove about 85% of color in culture containing 200 mg/L RB5 after 12 h. Complete decolorization was observed in less than 24 h. The dye concentration in control flasks remained unchanged. UV-vis spectra (200–800 nm) were studied. Reactive Black 5 presents two main absorption peaks, one in visible region (597 nm) and another in UV region (310 nm), which can be attributed to the presence of chromophoric azo bonds and both aryl and naphthalene-like moieties, respectively [2]. The results of our experiments showed that the peak at 597 nm disappeared approximately after 12 h (complete removal of this peak was observed in less than 24 h). Meanwhile the main peak at UV region (310 nm) was apparently shifted to 254 nm within the first 12 h and no more changes were observed during further incubation time up to 72 h.

3.1. Photocatalytic Experiments

Photocatalytic process was employed to investigate the degradation of RB5 at 50, 100, 200, 300 and 500 mg/L concentrations. Results (Figure 2) showed that this process was effective in the removal of RB5 at low concentrations; almost complete decolorization (94.8%) was achieved within 80 min of illumination time at 50 mg/L RB5. After 90 min of exposure time, dye decolorization at 100, 200, 300 and 500 mg/L RB5 concentrations was about 72.3%, 48.6%, 33.5% and 19.2%, respectively. The UV-vis absorptive spectra of RB5 (at 50 mg/L concentration) was studied at different exposure times in the photocatalytic experiments. Results (Figure 3) showed that absorbance in the visible and UV spectral regions decreased with increasing illumination time indicating the photocatalytic oxidation was effective in removal of two absorbance peaks in visible and UV regions which belong to azo bonds and aromatic rings (benzene and naphthalene), respectively.




Figure 2. Photodegradation of RB5 at different dye concentration




Figure 3. UV-visible spectra (200-800 nm) of RB5 (50 mg/L) at different time intervals

3.2. Decolorization Kinetics

Kinetics studies was performed for photocatalytic decolorization at 50, 100 and 200 mg/L RB5 concentrations and results (Figure 4) followed first-order kinetic model. This kinetic model was presented in equation of $-\ln C/C_0 = kt$, where C_0 , C , t and k are initial dye concentration, dye concentration at t (min), decolorization time (min) and decolorization rate constant (1/min), respectively [20]. Decolorization rate constant (k) was calculated from the slope of the curve. In Figure 4,

the plot of $\ln C/C_0$ versus t and the linear correlation between $-\ln C/C_0$ and illumination time are shown. R^2 and k values are presented in Table 1. Data showed that decolorization constant rate decreased as the initial dye concentration increased.

Table 1. Decolorization parameters in photocatalytic process at different concentrations of RB5

Dye concentration (mg/l)	R^{2*}	k^{\dagger} (1/min)
50	0.991	0.038
100	0.982	0.016
200	0.933	0.005

*: correlation coefficient.

†: decolorization rate constant.




Figure 4. The plot of $\ln C/C_0$ vs. time (at different concentrations of RB5) indicates that the kinetic studies follow first order reaction

3.3. Combined Biological-photochemical Process

As shown in Figure 5, at 200 mg/L dye concentration, 74.9% decolorization occurred after 4 h irradiation but the peak attributed to aromatic rings in UV region did not significantly change. Results (Figure 5, curves e,f,g) showed that application of biological-photocatalytic process was more efficient than the photocatalytic and biological process solely in aromatic byproduct remediation. Absorption changes were not significant between UV-vis spectra in combined process after 2 and 3 or 4 h illumination.




Figure 5. UV-vis spectral scans, a: Control (200 mg/L RB5 without treatment), b & c: after treatment by $TiO_2 + UV$ for 2 & 4 h, respectively, d: after treatment by *pseudomonas aeruginosa* ETL-2211 (24 h), e: after combined treatment by *pseudomonas aeruginosa* ETL-2211 (24 h) & $TiO_2 + UV$ (2 h), f& g: after combined processing by *pseudomonas aeruginosa* ETL-2211 (24 h) & $TiO_2 + UV$ (3 & 4 h respectively)

3.4. COD Measurements

COD gives a measure of dye degradation and generated intermediates during the irradiation as well as a measure of the oxygen equivalent of the organic content in a sample that is susceptible to oxidation by strong oxidant [17]. In the photocatalytic process, COD was measured in both, raw and treated dye, solutions. The COD for untreated sample (at 50 mg/L RB5 concentration) was about 25 mg/ L O_2 and for treated sample was not detectable presumably due to complete mineralization of the sample. In the combined process COD measurement was performed in three stages namely, before and after biological step and at 4 points in photocatalytic step (after treatment for 1, 2, 3 and 4 h). Biological treatment was able to remove 59.9% of COD. COD reduction was not observed during photochemical post treatment.

4. Discussion

Measures to develop environmentally favorable as well as cost-effective techniques to remove hazardous pollutants from textile wastewater, have received a great deal of attention. In the present study, application of biological, photocatalytic and combined processes was investigated in the treatment of synthetic medium containing RB5. In the photocatalytic process, the amount of decolorization decreased as RB5 concentration increased (Figure 2). At a high dye concentration, a significant amount of UV may be absorbed by the dye molecules rather than the TiO_2 particles and thus reduce the efficiency of the catalytic reaction. Another possible reason is the interference from intermediates formed upon degradation of the parental dye molecules. They may compete with the dye molecules for the limited adsorption and catalytic sites on the TiO_2 particles and thus inhibit decolorization. Such interference would be more pronounced in the presence of an elevated level of degradation intermediates formed upon an increased initial dye concentration [18,23]. COD measurement and UV-vis spectral analysis were used as a measure of dye degradation and generated intermediates during the processes. Mineralization of 50 mg/L RB5 solution was obtained after 80 min by photocatalytic process (in presence of 0.2 g/L TiO_2). COD value was not detectable after complete decolorization of 50 mg/L RB5 solution. However, photocatalytic process was not effective in the removal of the dye at high concentrations (≥ 200 mg/L). At 200 mg/L concentration, 74.9% of decolorization was achieved after 4 h illumination under photocatalytic process and the absorbance peak in UV region (attributed to aromatic rings) was not completely removed. These observations indicate the inadequacy of photocatalytic processes in removal of concentrated wastewater. Successful application of the biological treatment in decolorization of high concentrated synthetic medium (up to 1000 mg/L RB5) was noted but the remediation of aromatic rings produced from destruction of dye molecule did not occur (Figure 5), even with prolonged incubation time. AOP, followed by biological treatment, could be justified if bio-recalcitrant compounds are easily degradable by AOP and the resulting intermediates are easily degradable by the biological treatment [24], as

previously reported [4,23-29] in the degradation of biorecalcitrant organic compounds. A few reports describe the application of photocatalytic process as a post-treatment in a combined process [24,32,33]. These investigations were suggested that since the concentration of organic compounds were high in wastewater and the use of photocatalytic process as a pre-treatment stage was not favorable. González et al. [34] evaluated the degradation of chlorophenols by sequential biological-photochemical process using *Trametes pubescens* and TiO₂/UV. This process was suggested because partial oxidation of original molecules was occurred and intermediate molecules produced through biological process could be more toxic than original molecules [34]. In the present study, a two-step treatment process, namely, biological treatment by *Pseudomonas aeruginosa* ETL-2211 followed by TiO₂/UV photocatalytic degradation, was also assessed. In the combined process (at 200 mg/L RB), absorbance peak in UV region significantly disappeared after about 2 h of illumination (Figure 5) and about 60% of COD removal was achieved in the biological step but in photocatalytic step, COD reduction was not observed which could be due to the production of strongly oxidative by-products that are resistant to further oxidation by photocatalytic treatment. As previously reported by Tanaka et al. [35] further hydroxylation of aromatic intermediates products leads to the cleavage of the aromatic ring resulting in the formation of oxygen-containing aliphatic compounds [35]. It is apparent that the application of photocatalytic process for an extended period of time is not reasonable.

5. Conclusion

In the present study, COD measurement and UV-vis analysis were used as a measure of degradation of dye and generated intermediates during the processes. Successful application of the biological treatment in decolorization of high concentrated synthetic medium was noted; however not in the remediation of aromatic rings produced from destruction of dye molecule. Therefore, photocatalytic process was used as a post-treatment for degradation of aromatic rings and was effective for this purpose, but COD reduction was not observed in this process which could be due to the production of by-products that are resistant to further oxidation by photocatalytic treatment. Further evaluation is required to improve the system in which three stages take place. The third step (i.e., biological treatment) subsequent to the photocatalytic step could facilitate further COD reduction. Our results showed that the combined process was more effective than biological treatment in remediation of aromatic rings (resulting from decolorization of the dye) and was cost-effective (in terms of the electrical power utilized) compared to the photocatalytic treatment only.

References

- [1] Mantzavinos D, Psillakis E: Enhancement of biodegradability of industrial wastewaters by chemical oxidation pre-treatment: a review. *J Chem Technol Biotechnol* 2004, 79:431-454.
- [2] Lucas MS, Dias AA, Sampaio A, Amaral C, Peres JA: Degradação de um corante azo reativo por um processo combinado de tratamento biológico e fotocatalítico. *Water Res* 2007, 41:1103-1109.
- [3] Pratum C, Wongthanate J, Arunlertaree C, Prapagdee B: Decolorization of reactive dyes and textile dyeing effluent by *Pleurotus sajor-cajo*. *Int J Integr Biol* 2011, 11:52-57.
- [4] Sudarjanto G, Keller-Lehmann B, Keller J: Optimization of integrated chemical-biological degradation of a reactive azo dye using response surface methodology. *J Hazard Mater* 2006, B138:160-168.
- [5] Gregorio SD, Balestri F, Basile M, Matteini V, Gini F, Giansanti S, Tozzi MG, Basosi R, Lorenzi R: Sustainable discolorization of textile chromo-baths by spent mushroom substrate from the industrial cultivation of *Pleurotus ostreatus*. *J Environ Prot* 2010, 1:85-94.
- [6] Lucas MS, Amaral C, Sampaio A, Peres JA, Dias AA: Biodegradation of the diazo dye reactive black 5 by a wild isolate of *Candida oleophila*. *Enzyme Microb Tech* 2006, 39:51-55.
- [7] Chagas EP, Durrant LR: Decolorization of azo dyes by *Phanerochaete chrysosporium* and *Pleurotus sajorcajo*. *Enzyme Microb Tech* 2001, 29:473-477.
- [8] Banerjee P, De S: Treatment of textile effluent using advanced oxidation process and nanofiltration. *Int J Environ Eng* 2010, 2:56-72.
- [9] Aksu, Z.; Tezer, S. Equilibrium and kinetic modelling of biosorption of Remazol Black B by *Rhizopus arrhizus* in a batch system: effect of Temperature. *Process Biochemistry*.2000 (36) 431-439.
- [10] Kirby, N.; Marchant, R.; McMullan, G. Decolourisation of synthetic textile dyes by *Phlebia tremellosa*. *FEMS Microbiology Letters*. 2000 (188) 93-96.
- [11] Swamy, J.; Ramsay, J.A. The evaluation of white rot fungi in the decolouration of textile dyes. *Enz Microbial Technology*.1999 (24) 130-137.
- [12] Meehan, C.; Banat, IM.; McMullan, G.; Nigam, P.; Smyth, F.; Marchant, R. Decolorization of Remazol Black-B using a thermotolerant yeast, *Kluyveromyces marxianus* IMB3. *Environment Internatioanl*. 2000. (1-2) 75-79.
- [13] C. Vanderzant and D. F. Splittstoesser, *Compendium of Methods for the Microbiological Examination of Foods*, American public health association, Washington, DC, USA, 3rd edition, 1997.
- [14] M. Cheesbrough, *District Laboratory Practice in Tropical Countries*, Part 2, Cambridge University Press, London, UK, 2000.
- [15] J. G. Holt, N. R. Krieg, P. H. A. Sneath, J. T. Staley, and S. T. Williams, *Bergey's Manual of Determinative Bacteriology*, Lippincott Williams and Wilkins, Baltimore, Md, USA, 9th edition, 1994.
- [16] Jadhav SU, Kalme SD, Govindwar SP: Biodegradation of methyl red by *Galactomyces geotrichum* MTCC 1360. *Int Biodeter Biodegr* 2008, 62:135-142.
- [17] Mahmoodi NM, Arami M, Yousefi-Limaee N: Photocatalytic degradation of triazinic ring-containing azo dye (Reactive Red 198) by using immobilized TiO₂ photoreactor: Bench scale study. *J Hazard Mater* 2006, B133:113-118.
- [18] Karimi L, Zohoori S, Yazdanshenas ME: Photocatalytic degradation of azo dyes in aqueous solutions under UV irradiation using nano-strontium titanate as the nanophotocatalyst. *Soc: J Saudi Chem*; 2011.
- [19] Wong CL, Tan YN, Mohamed AR: Photocatalytic degradation of phenol using immobilized TiO₂ nanotube photocatalysts. *J Nanotech* 2011:1-9.
- [20] Mahvi AH, Ghanbarian M, Nasser S, Khairi A: Mineralization and discoloration of textile wastewater by TiO₂ nanoparticles. *Desalination* 2009, 239:309-316. 19.
- [21] APHA: Standard methods for the examination of water and wastewater. 18th edition. Washington, DC: American Public Health Association; 1992.
- [22] Bali U, Çatalkaya E, Şengül F: Photodegradation of reactive black 5, direct Red 28 and direct Yellow12 using UV, UV/H₂O₂ and UV/H₂O₂/Fe²⁺: a comparative study. *J Hazard Mater* 2004, B114:159-166.
- [23] So CM, Cheng MY, Yu JC, Wong PK: Degradation of azo dye Procion Red MX-5B by photocatalytic oxidation. *Chemosphere* 2002, 46:905-912.
- [24] Suryaman D, Hasegawa K, Kagaya S: Combined biological and photocatalytic treatment for the mineralization of phenol in water. *Chemosphere* 2006, 65:2502-2506.
- [25] Amat AM, Arques A, Beneyto H, García A, Miranda MA, Seguí S: Ozonisation coupled with biological degradation for treatment of

- phenolic pollutants: a mechanistically based study. *Chemosphere* 2003, 53:79-86.
- [26] Chan CY, Tao S, Dawson R, Wong PK: Treatment of atrazine by integrating photocatalytic and biological processes. *Environ Pollut* 2004, 131:45-54.
- [27] Feng F, Xu Z, Li X, You W, Zhen Y: Advanced treatment of dyeing wastewater towards reuse by the combined Fenton oxidation and membrane bioreactor process. *J Environ Sci* 2010, 22:1657-1665.
- [28] Genç N: Photocatalytic oxidation of a reactive azo dye and evaluation of the biodegradability of photocatalytically treated and untreated dye. *Water SA* 2004, 30(3):399-405.
- [29] Parra S, Malato S, Pulgarin C: New integrated photocatalytic-biological flow system using supported TiO₂ and fixed bacteria for the mineralization of isoproturon. *Appl Catal B –Environ* 2002, 36:131-144.
- [30] Tamer E, Aly AM, Ossama ET, Bo M, Benoit G: Sequential photochemicalbiological degradation of chlorophenols. *Chemosphere* 2007, 66:2201-2209.
- [31] Tamer E, Hamid Z, Aly AM, Ossama ET, Bo M, Benoit G: Sequential UVbiological degradation of chlorophenols. *Chemosphere* 2006, 63:277-284.
- [32] Balcioglu IA, Cecen F: Treatability of kraft pulp bleaching wastewater by biochemical and photocatalytic oxidation. *Wat Sci Technol* 1999, 40:281-288.
- [33] Li XY, Zhao YG: Advanced treatment of dyeing wastewater for reuse. *Wat Sci Technol* 1999, 39:249-255.
- [34] González LF, Sarria V, Sánchez OF: Degradation of chlorophenols by sequential biological-advanced oxidative process using *Trametes pubescens* and TiO₂/UV. *Bioresource Technol* 2010, 101:3493-3499.
- [35] Tanaka K, Robledo SM, Hisanaga T, Ali R, Ramli Z, Bakar WA: Photocatalytic degradation of 3,4-xylol N-methyl carbamate (MPMC) and other carbamate pesticides in aqueous TiO₂ suspensions. *J Mol Catal A-Chem* 1999, 144:425-430.