Application of Differentiation Term by Term Theorem on the Partial Differential Problems

Chii-Huei Yu*

Department of Management and Information, Nan Jeon University of Science and Technology, Tainan City, Taiwan *Corresponding author: chiihuei@nju.edu.tw

Received November 24, 2013; Revised January 05, 2014; Accepted January 16, 2014

Abstract This paper takes the mathematical software Maple as the auxiliary tool to study the partial differential problems of two types of two-variables functions. We can obtain the infinite series forms of any order partial derivatives of these two types of functions by using differentiation term by term theorem, and hence greatly reduce the difficulty of calculating their higher order partial derivative values. On the other hand, we propose two examples to do calculation practically. The research methods adopted in this study involved finding solutions through manual calculations and verifying our answers by using Maple.

Keywords: partial derivatives, infinite series forms, differentiation term by term theorem, Maple

Cite This Article: Chii-Huei Yu, "Application of Differentiation Term by Term Theorem on the Partial Differential Problems." *International Journal of Partial Differential Equations and Applications* 2, no. X (2014): 7-12. doi: 10.12691/ijpdea-2-1-2.

1. Introduction

information technology advances, computers can become comparable with human brains to perform abstract tasks, such as abstract art similar to the paintings of Picasso and musical compositions similar to those of Beethoven, is a natural question. Currently, this appears unattainable. In addition, whether computers can solve abstract and difficult mathematical problems and develop abstract mathematical theories such as those of mathematicians also appears unfeasible. Nevertheless, in seeking for alternatives, we can study what assistance mathematical software can provide. This study introduces how to conduct mathematical research using the mathematical software Maple. The main reasons of using Maple in this study are its simple instructions and ease of use, which enable beginners to learn the operating techniques in a short period. By employing the powerful computing capabilities of Maple, difficult problems can be easily solved. Even when Maple cannot determine the solution, problem-solving hints can be identified and inferred from the approximate values calculated and solutions to similar problems, as determined by Maple. For this reason, Maple can provide insights into scientific research.

In calculus and engineering mathematics curricula, the evaluation and numerical calculation of the partial derivatives of multivariable functions are important. For example, Laplace equation, wave equation, as well as other important physical equations are involved the partial derivatives. On the other hand, calculating the q-th order partial derivative value of a multivariable function at some point, in general, needs to go through two procedures:

firstly determining the q-th order partial derivative of this function, and then taking the point into the q-th order partial derivative. These two procedures will make us face with increasingly complex calculations when calculating higher order partial derivative values (i.e. q is large), and hence to obtain the answers by manual calculations is not easy. In this article, we study the partial differential problem of the following two types of two-variables functions

$$f(x,y) = \frac{\left(\sqrt{\alpha^2 x^2 + \beta^2 y^2}\right)^k}{\sqrt{(1+\alpha x)^2 + \beta^2 y^2}} \cos\left[k \cot^{-1}\left(\frac{\alpha x}{\beta y}\right) - \cot^{-1}\left(\frac{1+\alpha x}{\beta y}\right)\right]$$
(1)

$$g(x,y) = \frac{\left(\sqrt{\alpha^2 x^2 + \beta^2 y^2}\right)^k}{\sqrt{(1+\alpha x)^2 + \beta^2 y^2}} \sin\left[k \cot^{-1}\left(\frac{\alpha x}{\beta y}\right) - \cot^{-1}\left(\frac{1+\alpha x}{\beta y}\right)\right]$$
(2)

where α, β are real numbers, $\beta \neq 0$, and k is any integer. We can obtain the infinite series forms of any order partial derivatives of these two types of two-variables functions using differentiation term by term theorem; these are the major results of this study (i.e., Theorems 1 and 2), and hence greatly reduce the difficulty of calculating their higher order partial derivative values. As for the study of related partial differential problems can refer to [1-13]. In addition, we provide two examples to do calculation practically. The research methods adopted in this study involved finding solutions through manual calculations and verifying these solutions by using Maple. This type of research method not only allows the discovery of calculation errors, but also helps modify the original

directions of thinking from manual and Maple calculations. Therefore, Maple provides insights and guidance regarding problem-solving methods.

2. Main Results

Firstly, we introduce some notations and formulas used in this study.

2.1. Notations

- **2.1.1.** Let z = a + ib be a complex number, where $i = \sqrt{-1}$, a, b are real numbers. We denote a the real part of z by Re(z), and b the imaginary part of z by Im(z).
- **2.1.2.** Suppose m, n are non-negative integers. For the two-variables function f(x, y), its n-times partial derivative with respect to x, and m-times partial derivative with respect to y, forms a m+n-th order partial derivative,

and denoted by
$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)$$
.

2.1.3. Suppose r is any real number, m is any positive integer. Define $(r)_m = r(r-1)...(r-m+1)$, and $(r)_0 = 1$.

2.2. Formulas

2.2.1. Euler's Formula

 $e^{i\theta} = \cos \theta + i \sin \theta$, where θ is any real number.

2.2.2. DeMoivre's Formula

 $(\cos \theta + i \sin \theta)^k = \cos k\theta + i \sin k\theta$, where k is any integer, and θ is any real number.

Next, we introduce an important theorem used in this paper.

2.3. Differentiation Term by Term Theorem ([14])

For all non-negative integers k, if the functions $g_k:(a,b)\to R$ satisfy the following three conditions : (i)

there exists a point $x_0 \in (a,b)$ such that $\sum_{k=0}^{\infty} g_k(x_0)$ is

convergent, (ii) all functions $g_k(x)$ are differentiable on

open interval (a,b), (iii) $\sum_{k=0}^{\infty} \frac{d}{dx} g_k(x)$ is uniformly

convergent on (a,b) . Then $\sum_{k=0}^{\infty} g_k(x)$ is uniformly

convergent and differentiable on (a,b). Moreover, its

derivative
$$\frac{d}{dx} \sum_{k=0}^{\infty} g_k(x) = \sum_{k=0}^{\infty} \frac{d}{dx} g_k(x)$$
.

Before deriving the first major result in this study, we need two lemmas.

2.4. Lemma 1

Suppose z is a complex number with $|z| \neq 1$ and k is any integer. Then

$$\frac{z^{k}}{1+z} = \begin{cases}
\sum_{p=0}^{\infty} (-1)^{p} z^{p+k} & \text{if } |z| < 1 \\
\sum_{p=0}^{\infty} (-1)^{p} z^{-p+k-1} & \text{if } |z| > 1
\end{cases} \tag{3}$$

Proof If |z| < 1, then

$$\frac{z^k}{1+z} = z^k \sum_{p=0}^{\infty} (-1)^p z^p$$
$$= \sum_{p=0}^{\infty} (-1)^p z^{p+k}$$

If |z| > 1, then

$$\frac{z^k}{1+z} = \frac{z^{k-1}}{1+1/z}$$

$$= z^{k-1} \sum_{p=0}^{\infty} (-1)^p z^{-p}$$

$$= \sum_{p=0}^{\infty} (-1)^p z^{-p+k-1}$$

2.5. Lemma 2

Suppose a,b are real numbers, b>0, and k is an integer. Then

$$(a+ib)^k = (a^2+b^2)^{k/2} \exp\left[ik \cot^{-1}\frac{a}{b}\right]$$
 (5)

Proof $(a+ib)^k$

$$= \left[\sqrt{a^2 + b^2} \left(\frac{a}{\sqrt{a^2 + b^2}} + i\frac{b}{\sqrt{a^2 + b^2}}\right)\right]^k$$
$$= (a^2 + b^2)^{k/2} \cdot (\cos\theta + i\sin\theta)^k$$

(where
$$\theta = \cot^{-1} \frac{a}{b}$$
)
= $(a^2 + b^2)^{k/2} \cdot (\cos k\theta + i \sin k\theta)$

(By DeMoivre's formula)

$$= (a^2 + b^2)^{k/2} \exp\left[ik \cot^{-1}\frac{a}{b}\right]$$

(By Euler's formula)

Next, we determine the infinite series forms of any order partial derivatives of the two-variables function (1).

2.6. Theorem 1

Suppose α, β are real numbers, $\beta \neq 0$, m, n are non-negative integers, and k is an integer. If the domain of two-variables function

$$f(x,y) = \frac{\left(\sqrt{\alpha^2 x^2 + \beta^2 y^2}\right)^k}{\sqrt{(1+\alpha x)^2 + \beta^2 y^2}} \cos\left[k \cot^{-1}\left(\frac{\alpha x}{\beta y}\right) - \cot^{-1}\left(\frac{1+\alpha x}{\beta y}\right)\right]$$

is
$$\{(x, y) \in \mathbb{R}^2 \mid \beta y > 0, \alpha^2 x^2 + \beta^2 y^2 \neq 1 \}$$
.

Case (1) If $\alpha^2 x^2 + \beta^2 y^2 < 1$ and $\beta y > 0$. Then the m+n-th order partial derivative of f(x,y),

$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{p+k-m-n} \times (6)$$

$$\cos \left[(p+k-m-n) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$

Case (2) If $\alpha^2 x^2 + \beta^2 y^2 > 1$ and $\beta y > 0$. Then

$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (-p+k-1)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{-p+k-m-n-1} \times \cos \left[(-p+k-m-n-1) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$
(7)

Proof Let $z = \alpha x + i\beta y$, then

$$\frac{\frac{z^k}{1+z}}{\frac{1+z}{1+\alpha x+i\beta y}} = \frac{(\alpha x+i\beta y)^k}{1+\alpha x+i\beta y} = \frac{\left(\sqrt{\alpha^2 x^2+\beta^2 y^2}\right)^k \exp\left[ik\cot^{-1}\frac{\alpha x}{\beta y}\right]}{\sqrt{(1+\alpha x)^2+\beta^2 y^2} \exp\left[i\cot^{-1}\left(\frac{1+\alpha x}{\beta y}\right)\right]}$$

(By Lemma 2)

$$= \frac{\left(\sqrt{\alpha^2 x^2 + \beta^2 y^2}\right)^k}{\sqrt{(1 + \alpha x)^2 + \beta^2 y^2}} \times \exp\left[i\left(k\cot^{-1}\frac{\alpha x}{\beta y} - \cot^{-1}\frac{1 + \alpha x}{\beta y}\right)\right] (8)$$

Therefore,

$$f(x, y) = \operatorname{Re}\left(\frac{z^k}{1+z}\right) \tag{9}$$

Case (1) If $\alpha^2 x^2 + \beta^2 y^2 < 1$ and $\beta y > 0$. By (3) and (9), we obtain

$$f(x, y) = \operatorname{Re}\left(\sum_{p=0}^{\infty} (-1)^{p} (\alpha x + i\beta y)^{p+k}\right)$$
 (10)

Using differentiation term by term theorem, differentiating n-times with respect to x, and m-times with respect to y on both sides of (10), we have

$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)$$

$$= \operatorname{Re} \left(\alpha^n (i\beta)^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} (\alpha x + i\beta y)^{p+k-m-n} \right)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} \operatorname{Re} [i^m (\alpha x + i\beta y)^{p+k-m-n}]$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{p+k-m-n} \times$$

$$\operatorname{Re} \left[\exp \left(i \frac{m\pi}{2} \right) \exp \left[i (p+k-m-n) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) \right] \right]$$
(By Lemma 2)
$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{p+k-m-n} \times$$

$$\cos \left[(p+k-m-n) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$

Case (2) If $\alpha^2 x^2 + \beta^2 y^2 > 1$ and $\beta y > 0$. Using (4) and (9), we have

$$f(x,y) = \text{Re}\left(\sum_{p=0}^{\infty} (-1)^p (\alpha x + i\beta y)^{-p+k-1}\right)$$
 (11)

Also, by differentiation term by term theorem, we obtain

$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)$$

$$= \operatorname{Re} \left(\alpha^n (i\beta)^m \sum_{p=0}^{\infty} (-1)^p (-p+k-1)_{m+n} (\alpha x + i\beta y)^{-p+k-m-n-1} \right)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (-p+k-1)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{-p+k-m-n-1} \times$$

$$\operatorname{Re} \left[\exp \left(i \frac{m\pi}{2} \right) \exp \left[i (-p+k-m-n-1) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) \right] \right]$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (-p+k-1)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{-p+k-m-n-1} \times$$

$$\operatorname{cos} \left[(-p+k-m-n-1) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$

Next, we find the infinite series forms of any order partial derivatives of the two-variables function (2).

2.7. Theorem 2

If the assumptions are the same as Theorem 1. Suppose the domain of the two-variables function

$$g(x,y) = \frac{\left(\sqrt{\alpha^2 x^2 + \beta^2 y^2}\right)^k}{\sqrt{(1+\alpha x)^2 + \beta^2 y^2}} \sin\left[k \cot^{-1}\left(\frac{\alpha x}{\beta y}\right) - \cot^{-1}\left(\frac{1+\alpha x}{\beta y}\right)\right]$$

is
$$\{(x, y) \in \mathbb{R}^2 \mid \beta y > 0, \alpha^2 x^2 + \beta^2 y^2 \neq 1 \}$$
.

Case (1) If $\alpha^2 x^2 + \beta^2 y^2 < 1$ and $\beta y > 0$. Then the m+n-th order partial derivative of g(x,y),

$$\frac{\partial^{m+n} g}{\partial y^m \partial x^n}(x, y)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{p+k-m-n} \times \sin \left[(p+k-m-n) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$
(12)

Case (2) If $\alpha^2 x^2 + \beta^2 y^2 > 1$ and $\beta y > 0$. Then

$$\frac{\partial^{m+n} g}{\partial y^m \partial x^n}(x, y)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (-p+k-1)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{-p+k-m-n-1} \times \sin \left[(-p+k-m-n-1)\cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$
(13)

Proof By (8), we obtain

$$g(x, y) = \operatorname{Im}\left(\frac{z^k}{1+z}\right) \tag{14}$$

Case (1) If $\alpha^2 x^2 + \beta^2 y^2 < 1$ and $\beta y > 0$. Using (3) and (14), we have

$$g(x,y) = \operatorname{Im}\left(\sum_{p=0}^{\infty} (-1)^p (\alpha x + i\beta y)^{p+k}\right)$$
(15)

Thus, by differentiation term by term theorem,

$$\frac{\partial^{m+n} g}{\partial y^m \partial x^n}(x, y)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (p+k)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{p+k-m-n} \times \sin \left[(p+k-m-n) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$

Case (2) If $\alpha^2 x^2 + \beta^2 y^2 > 1$ and $\beta y > 0$. Then using (4) and (14), we obtain

$$g(x, y) = \text{Im} \left(\sum_{p=0}^{\infty} (-1)^p (\alpha x + i\beta y)^{-p+k-1} \right)$$
 (16)

Also, by differentiation term by term theorem,

$$\frac{\partial^{m+n} g}{\partial y^m \partial x^n}(x, y)$$

$$= \alpha^n \beta^m \sum_{p=0}^{\infty} (-1)^p (-p+k-1)_{m+n} \left(\sqrt{\alpha^2 x^2 + \beta^2 y^2} \right)^{-p+k-m-n-1} \times \sin \left[(-p+k-m-n-1) \cot^{-1} \left(\frac{\alpha x}{\beta y} \right) + \frac{m\pi}{2} \right]$$

3. Examples

In the following, for the partial differential problem of the two types of two-variables functions in this study, we propose two examples and use Theorems 1, 2 to determine the infinite series forms of any order partial derivatives of these functions, and evaluate some of their higher order partial derivative values. On the other hand, we employ Maple to calculate the approximations of these higher order partial derivative values and their solutions for verifying our answers.

$$f(x,y) = \frac{\left(\sqrt{4x^2 + 9y^2}\right)^5}{\sqrt{(1+2x)^2 + 9y^2}} \cos \begin{bmatrix} 5\cot^{-1}\left(\frac{2x}{3y}\right) \\ -\cot^{-1}\left(\frac{1+2x}{3y}\right) \end{bmatrix}$$
(17)

is
$$\{(x, y) \in \mathbb{R}^2 \mid y > 0, 4x^2 + 9y^2 \neq 1 \}$$
.

Case (1) If $4x^2 + 9y^2 < 1$ and y > 0. Then by (6), we obtain any m+n-th order partial derivative of f(x, y),

$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)$$

$$= 2^n 3^m \sum_{p=0}^{\infty} (-1)^p (p+5)_{m+n} \left(\sqrt{4x^2 + 9y^2}\right)^{p+5-m-n} \times (18)$$

$$\cos \left[(p+5-m-n)\cot^{-1} \left(\frac{2x}{3y}\right) + \frac{m\pi}{2} \right]$$

Thus, we can evaluate the 13-th order partial derivative value of f(x, y) at $\left(\frac{1}{4}, \frac{1}{9}\right)$,

$$\frac{\partial^{13} f}{\partial y^{6} \partial x^{7}} \left(\frac{1}{4}, \frac{1}{9} \right)$$

$$= -2^{7} 3^{6} \sum_{p=0}^{\infty} (-1)^{p} (p+5)_{13} \left(\frac{\sqrt{13}}{6} \right)^{p-8} \cos \left[(p-8) \cot^{-1} \left(\frac{3}{2} \right) \right]$$
(19)

Next, we use Maple to verify the correctness of (19). >f:=(x,y)->(sqrt(4*x^2+9*y^2))^5/sqrt((1+2*x)^2+9*y^2)*cos(5*arccot((2*x)/(3*y))-arccot((1+2*x)/(3*y))); >evalf(D[1\$7,2\$6](f)(1/4,1/9),18);

$$1.41580578711082968 \cdot 10^{12}$$

>evalf(-2^7*3^6*sum((-1)^p*product(p+5-j,j=0..12)*(sqrt (13)/6)^(p-8)*cos((p-8)*arccot(3/2)),p=0..infinity),18);

$$1.41580578711082969 \cdot 10^{12}$$

Case (2) If $4x^2 + 9y^2 > 1$ and y > 0. Using (7), we obtain

$$\frac{\partial^{m+n} f}{\partial y^m \partial x^n}(x, y)
= 2^n 3^m \sum_{p=0}^{\infty} (-1)^p (-p+4)_{m+n} \left(\sqrt{4x^2 + 9y^2}\right)^{-p+4-m-n}
\times \cos\left[\left(-p+4-m-n\right)\cot^{-1}\left(\frac{2x}{3y}\right) + \frac{m\pi}{2}\right]$$
(20)

Hence.

$$\frac{\partial^{9} f}{\partial y^{4} \partial x^{5}} (2,7)$$

$$= 2^{5} 3^{4} \sum_{p=0}^{\infty} (-1)^{p} (-p+4)_{9} \left(\sqrt{457}\right)^{-p-5} \cos \left[(-p-5) \cot^{-1} \left(\frac{4}{21}\right) \right]$$
(21)

>evalf(D[1\$5,2\$4](f)(2,7),18);

0.0000296927268090019832

>evalf(2^5*3^4*sum((-1)^p*product(-p+4-j,j=0..8)*(sqrt (457))^(-p-5)*cos((-p-5)*arccot(4/21)),p=0..infinity),18); 0.0000296927268090019828

3.2. Example 2

If the domain of the two-variables function

$$g(x,y) = \frac{\left(\sqrt{16x^2 + y^2}\right)^7}{\sqrt{(1+4x)^2 + y^2}} \sin\left[7\cot^{-1}\left(\frac{4x}{y}\right) - \cot^{-1}\left(\frac{1+4x}{y}\right)\right]$$
(22)

is
$$\{(x, y) \in \mathbb{R}^2 \mid y > 0, 16x^2 + y^2 \neq 1 \}$$
.

Case (1) If $16x^2 + y^2 < 1$ and y > 0. By (12), we obtain any m+n-th order partial derivative of g(x, y),

$$\frac{\partial^{m+n} g}{\partial y^m \partial x^n}(x, y)$$

$$= 4^n \sum_{p=0}^{\infty} (-1)^p (p+7)_{m+n} \left(\sqrt{16x^2 + y^2} \right)^{p+7-m-n} \times (23)^n$$

$$\sin \left[(p+7-m-n)\cot^{-1} \left(\frac{4x}{y} \right) + \frac{m\pi}{2} \right]$$

It follows that

$$\frac{\partial^{11}g}{\partial y^{4}\partial x^{7}} \left(\frac{1}{6}, \frac{1}{2}\right)$$

$$= 4^{7} \sum_{p=0}^{\infty} (-1)^{p} (p+7)_{11} \left(\frac{5}{6}\right)^{p-4} \sin\left[(p-4)\cot^{-1}\left(\frac{4}{3}\right)\right]$$
(24)

>g:=(x,y)-> $(sqrt(16*x^2+y^2))^7/sqrt((1+4*x)^2+y^2)*sin(7*arccot((4*x)/y)-arccot((1+4*x)/y));$ >evalf(D[1\$7,2\$4](g)(1/6,1/2),18);

 $2.95759934660351636 \cdot 10^{8}$

>evalf(4^7*sum((-1)^p*product(p+7-j,j=0..10)*(5/6)^(p-4)*sin((p-4)*arccot(4/3)),p=0..infinity),18);

$$2.95759934660351647 \cdot 10^{8}$$

Case (2) If
$$16x^2 + y^2 > 1$$
 and $y > 0$. Using (13), we have
$$\frac{\partial^{m+n} g}{\partial y^m \partial x^n}(x, y)$$

$$=4^{n} \sum_{p=0}^{\infty} (-1)^{p} (-p+6)_{m+n} \left(\sqrt{16x^{2} + y^{2}} \right)^{-p+6-m-n} \times (25)$$

$$\sin \left[(-p+6-m-n) \cot^{-1} \left(\frac{4x}{y} \right) + \frac{m\pi}{2} \right]$$

Therefore,

$$\frac{\partial^{12} g}{\partial y^{6} \partial x^{6}} (3,4)$$

$$= -4^{6} \sum_{p=0}^{\infty} (-1)^{p} (-p+6)_{12} (\sqrt{160})^{-p-6} \sin \left[(-p-6) \cot^{-1} 3 \right]$$
(26)

>evalf(D[1\$6,2\$6](g)(3,4),18);

0.00242325523123641135

>evalf(-4^6*sum((-1)^p*product(-p+6-j,j=0..11)*(sqrt(160))^(-p-6)*sin((-p-6)*arccot(3)),p=0..infinity),18);

0.00242325523123641134

4. Conclusion

In this article, we provide a new technique to evaluate any order partial derivatives of two types of two-variables functions. We hope this technique can be applied to solve another partial differential problems. On the other hand, the differentiation term by term theorem plays a significant role in the theoretical inferences of this study. In fact, the applications of this theorem are extensive, and can be used to easily solve many difficult problems; we endeavor to conduct further studies on related applications. In addition, Maple also plays a vital assistive role in problem-solving. In the future, we will extend the research topic to other calculus and engineering mathematics problems and solve these problems by using Maple. These results will be used as teaching materials for Maple on education and research to enhance the connotations of calculus and engineering mathematics.

References

- L. E. Fraenkel, "Formulae for high derivatives of composite functions," Mathematical Proceedings of the Cambridge Philosophical Society, Vol. 83, pp. 159-165, 1978.
- [2] T-W, Ma, "Higher chain formula proved by combinatorics," The Electronic Journal of Combinatorics, Vol. 16, #N21, 2009.
- [3] D. N. Richard, "An efficient method for the numerical evaluation of partial derivatives of arbitrary order," ACM Transactions on Mathematical Software, Vol. 18, No. 2, pp. 159-173, 1992.
- [4] C. H., Bischof, G. Corliss, and A. Griewank, "Structured second and higher-order derivatives through univariate Taylor series," *Optimization Methods and Software*, Vol. 2, pp. 211-232, 1993.
- [5] A. Griewank and A. Walther, Evaluating derivatives: principles and techniques of algorithmic differentiation, 2nd ed., SIAM, Philadelphia, 2008.
- [6] C.-H. Yu, "Using Maple to evaluate the partial derivatives of twovariables functions," *International Journal of Computer Science* and Mobile Computing, Vol. 2, Issue. 6, pp. 225-232, 2013.
- [7] C.-H. Yu, "Using Maple to study the partial differential problems," Applied Mechanics and Materials, in press, 2013.
- [8] C. -H. Yu, "Evaluating partial derivatives of two-variables functions by using Maple," Proceedings of the 6th IEEE/International Conference on Advanced Infocomm Technology, Taiwan, pp. 23-27, 2013.
- [9] C.-H. Yu, "Application of Maple: taking the partial differential problem of some types of two-variables functions as an example," *Proceedings of the International Conference on e-Learning*, Taiwan, pp. 337-345, 2013.
- [10] C.-H. Yu, "Application of Maple on the partial differential problem of four types of two-variables functions," Proceedings of the International Conference on Advanced Information Technologies, Taiwan, No. 87, 2013.

- [11] C.-H., Yu, "Application of Maple: taking the partial differential problem of two-variables functions as an example," *Proceedings* of 2013 Business Innovation and Development Symposium, Taiwan, B20130113001, 2013.
- [12] C. -H. Yu and B. -H. Chen, "The partial differential problem," *Computational Research*, Vol. 1, No. 3, pp. 53-60, 2013.
- [13] C. -H. Yu, "Partial derivatives of some types of two-variables functions," *Pure and Applied Mathematics Journal*, Vol. 2, No. 2, pp. 56-61, 2013.
- [14] T. M. Apostol, *Mathematical analysis*, 2nd ed., Addison-Wesley, Boston, p 230, 1975.