

Airborne Lead in El Paso, Texas, USA

Paz L.M.¹, Amaya M.A.², Clague J.W.³, Li W-W⁴, Olvera H.A.², Berwick M.⁵, Burchiel S.W.⁶, Pingitore N.E.^{1,*}

¹Department of Geological Sciences, The University of Texas at El Paso, El Paso, Texas, USA

²School of Nursing, The University of Texas at El Paso, El Paso, Texas, USA

³Environmental Engineering & Public Health (E2PH), LLC, El Paso, Texas, USA

⁴Department of Civil Engineering, The University of Texas at El Paso, El Paso, Texas, USA

⁵Department of Internal Medicine and Dermatology, The University of New Mexico, Albuquerque, NM, USA

⁶College of Pharmacy, The University of New Mexico, Albuquerque, NM, USA

*Corresponding author: npingitore@utep.edu

Abstract Despite significant strides to minimize lead in our air, water, soil, food, homes, workplaces, and consumer products, sporadic and systemic lead poisoning persists in industrial societies, e.g., the contemporary Flint (Michigan, USA) municipal water tragedy. Because lead exposure is cumulative, the sum of exposure from all sources, it is important to document and update potential community or neighborhood exposure levels from such environmental compartments as air, water, soil, food, and housing. Here we present ambient airborne lead levels in El Paso, Texas, USA, parsed by particulate matter (PM) size fraction, geography, and season. Dichotomous samplers at 8 stations collected PM continuously for 1-week periods for nearly 4 years. Overall, airborne annual lead exposure throughout El Paso County is low, well within US Environmental Protection Agency (EPA) guidelines. There is considerably more lead in the coarse fraction (PM₁₀-PM_{2.5}) than in the fine fraction (PM_{2.5} and lower), indicating decreased effective human exposures because the coarse fraction does not penetrate as deeply into the pulmonary system as finer particles. Re-entrained soil has previously been identified as the source of airborne lead in El Paso; the concentration of lead in the coarse PM is consistent with this observation. Seasonally, fall-winter lead levels are highest, due to PM trapping by temperature inversions, followed by spring and summer. Geospatially, higher lead levels characterize sampling stations in older and more commercial neighborhoods, also consistent with published maps of soil lead levels throughout the city and county. At present, airborne lead can be considered to be only a minor source of lead exposure for typical El Paso residents.

Keywords: lead, air pollution, particulate matter, El Paso

Cite This Article: Paz L.M., Amaya M.A., Clague J.W., Li W-W, Olvera H.A., Berwick M., Burchiel S.W., and Pingitore N.E., "Airborne Lead in El Paso, Texas, USA." *Journal of Atmospheric Pollution*, vol. 5, no. 2 (2017): 47-54. doi: 10.12691/jap-5-2-2.

1. Introduction

Airborne lead (Pb) contamination was a serious environmental health hazard in industrialized countries through much of the 20th Century [1,2,3,4]. Leaded gasoline was the dominant source, rivalled only in specific local areas by fugitive emissions from mining, smelting, and working of this toxic metal, as well as incineration and such specific industries as lead-acid battery and wheel weight manufacture. The addition of tetra-ethyl lead and related Pb-based anti-knock compounds decreased in the US as catalytic converters, which would be inactivated by lead deposition, were incorporated into new vehicles to reduce smog, starting in 1975 [5]. A consequent dramatic decrease in blood lead levels correlated closely with the measured decrease in airborne lead over the next 15 or so years [6]. Similar decreases were observed in other nations that switched to unleaded gasoline, a process that was completed in most industrialized nations by 2000 and globally by 2015.

The most dangerous aspect of human exposure to lead occurs in young children, in whom lead acts as a powerful neurotoxin on the developing brain and nerve system, even at quite low blood lead levels (BLL) [7,8,9]. Current practice in the US, as enunciated by The Centers for Disease Control and Prevention (CDC), employs a blood lead intervention threshold in children of ≥ 10 $\mu\text{g}/\text{dL}$. In the US average BLL in children 1-5 years old has dropped from 15 to less than 1.5 $\mu\text{g}/\text{dL}$ [10]. Nonetheless, some researchers and activists argue that no levels of blood lead are safe [11]. In 2008 the US Environmental Protection Agency (EPA) lowered the then current airborne lead standard by a factor of 10, from 1.5 $\mu\text{g m}^{-3}$ to a 3-month average below 0.15 $\mu\text{g m}^{-3}$ in total suspended particulate matter (TSP) [12,13]. The agency also relaxed the required monitoring for urban areas since most were expected to meet the new, albeit stricter, regulation. Nonetheless, with a decrease in the number of required urban monitors it is possible that in some cities there may be persistent local "hot spots" that do not meet current regulations.

1.1. Legacy of Lead Contamination in El Paso, Texas

The city of El Paso, in far west Texas, USA, has a long history of exposure to lead due in part to the presence for more than a century of a lead smelter sited approximately 3 km from the urban downtown center. The smelter was constructed in 1887 to process ores from mines in Mexico and the US Southwest. Over time it was chiefly a lead smelter and then a copper smelter. Operations were suspended in 1999 and the smelter was demolished in 2013. The soil around the smelter was contaminated with lead and arsenic, and children in Smelertown, a workers' residential community immediately adjacent to the property, were found to have exceedingly high BLLs [14]. The children were treated and the settlement was razed. Extensive background material on Smelertown and the ASARCO smelter are found in [15,16].

With cessation of smelter operations and elimination of leaded gasoline, airborne lead in El Paso has fallen dramatically [17]. Nonetheless, Pb-contaminated soils persist in the older areas of El Paso, a legacy of smelting, gasoline, and other industrial activities [18,19]. Selected residential yards in a small region near the smelter site were remediated under EPA supervision.

By speciating recent samples of airborne lead by synchrotron-based x-ray absorption spectroscopy, re-entrained soil was found to be the dominant current source of the metal in the El Paso air shed [20].

1.2. El Paso Geography

El Paso is located in the far west region of Texas, along the Rio Grande, which forms that section of the USA-Mexico border (Figure 1). El Paso County hosts some 825,000 residents, with an estimated 1,500,000 residents in contiguous Cd. Juarez, Chihuahua, Mexico. The climate is Chihuahuan Desert, a high desert climate zone characterized by low rainfall and cool to cold winter temperatures [21]. El Paso elevation is approximately 1150 m, with an average rainfall of just under 25 cm, most of which occurs in brief mid-to-late-summer storms in the "monsoon" season of July, August, and September. Strong winds and dust storms characterize the dry spring season.

1.3. Rationale for Study

This study presents the results of lead analyses of a large number of dichotomous aerosol samples, which

split particulate matter (PM) samples into PM coarse ($PM_{10} - PM_{2.5}$) and PM fine ($PM_{2.5}$ and finer). Because sampling was performed at numerous sites throughout the city of El Paso for periods of several years, the sample set can provide a robust picture of contemporary exposure of the local citizenry to airborne lead.



Figure 1. (Color online) Location map of El Paso, Texas

2. Materials and Methods

2.1. Sampling

The samples included in this study were collected at 8 stations over varied periods of time during the 3.4 years from August 2006 to December 2009. Several additional stations with shorter sampling periods were not included here. The sampling sites were widely dispersed across El Paso County, as seen in Figure 2. Sampling locations required access to electric power and security. Therefore we co-located most of our sites with existing Texas Commission on Environmental Quality (TCEQ, the Texas State agency responsible to the federal EPA) Community Air Monitoring Stations (CAMS) [22]. In areas without such CAMS, samplers were sited at El Paso Water Utilities (EPWU) ground water monitoring stations. A map of the sites is seen in Figure 2; brown shading is undeveloped mountain areas (Franklin Mts., USA and Sierra de Juarez, Mexico).

Descriptions of the individual sites are seen in Table 1. Sites B, C, E, and G were not included in this study.

Table 1. Description of 8 monitoring sites in El Paso, Texas

Site	Description
A	Predominantly residential area in west El Paso, adjacent to railroad tracks.
D	Northeast El Paso, classified as a residential location.
F	Adjacent to the campus of the University of Texas at El Paso (UTEP). Residential and some commercial.
H	Adjacent to the International Bridge of the Americas, a major US—Mexico border crossing and truck route.
I	Predominantly residential area in east El Paso, near Album Park.
J	Far southwest area of El Paso, the mixed urban—rural Mission Valley.
K	Southwest area of El Paso (Mission Valley), chiefly residential.
L	Monte Vista, newer residential area in easternmost El Paso, proximal to desert.




Figure 2. (Color online) Sample locations in El Paso County

2.2. Samples

Thermo Electron Corporation Series 241 dichotomous samplers (Thermo Electron Corp., Waltham, MA, USA) collected PM samples on Teflon® filters over a sampling period of 1 week. Mass was determined by gravimetric analysis. U.S. EPA's Compendium Method IO-2.2 guided overall procedures [23], with specifics available in [24].

2.3. Lead Determination

Lead concentration was determined on a PANalytical Epsilon5 energy dispersive x-ray fluorescence unit (XRF), an instrument specifically designed to meet the requirements of US EPA Method IO-3.3 for elemental analysis of air filters [25]. NIST PM SRMs (US National Institute for Standards and Technology, Standard Reference Materials) for particulate matter [26] and Micromatter™ commercial thin-film single-element standards [27] were used for instrument calibration. Analytical protocol was based on EPA Method IO-3.3 for XRF analysis of suspended particulate matter [25].

2.4. Data Compilation: Seasonality and Geography

To assist in comparisons and interpretation, the weekly data were composited into 3 annual seasons, defined by changing wind patterns. For the purposes of this research, June 16 through October 15 represented the 'summer' season, which is most commonly associated with high temperatures and intermittent brief monsoon rains (the North American Monsoon, bringing moisture to the region from the Gulf of Mexico). October 16 through February

14 data were designated as 'winter,' with area inversions typical for this part of the year. Nighttime freezing temperatures are not uncommon in this Chihuahuan Desert climate, at an elevation of ~1200 m. February 15 through June 15 was categorized as 'spring,' which is characterized by dry conditions and large-scale wind events. Each season has distinct characteristics associated with prevailing wind direction, temperature, and presence or absence of precipitation. Thus weekly data were simplified by calculating the average mass of lead per cubic meter of air, in each season, at each of the 8 sites, as averaged during a period of approximately four years.

The averages, organized based on season and geographic location, represent the total ambient exposure of a person residing in a specific El Paso neighborhood during a specific time period. This represents an important parameter in considering the health impact of different neighborhoods on their residents.

3. Results and Discussion

3.1. Data

Figure 3 presents the seasonal averages of lead in PM_C (PM_{coarse}) for all stations for all years. Figure 4 presents the seasonal averages of lead in PM_F (PM_{fine}) for all stations for all years. Note the change in vertical scales, where the PM_C tops out at 60 ng m^{-3} and PM_F tops out at only 30 ng m^{-3} . Thus differences between lead levels in coarse and fine PM are twice as great as they appear visually. Had we plotted the PM_F at the same scale as the PM_C , differences *within* the PM_F data set would not have been visible.




Figure 3. (Color online) PM_c lead levels by season and station




Figure 4. (Color online) PM_f lead levels by season and station




Figure 5. (Color online) Comparison of lead in total suspended particulates for cities located in the southwest U.S.

3.2. General Observations

3.2.1. Lead Levels and Air Quality Regulations

Airborne lead was found to be well below the US EPA National Ambient Air Quality Standards (NAAQS) for Lead (Pb) under the Clean Air Act of 0.15 micrograms Pb in total suspended particles per cubic meter air as a 3-month average. The El Paso Pb values are seen to have not exceeded an average of approximately 70 ng m^{-3} , or $0.07 \mu\text{g m}^{-3}$, during any of our averaged periods of approximately 4 months. Note that we used the sum of our PM_{10} and $\text{PM}_{2.5}$ measurements to derive 70 ng m^{-3} as a proxy for the TSP value used in the regulations.

Figure 5 presents a comparison of airborne lead levels in El Paso and 3 other cities in the region, as well as the US average and US EPA standard [28,29]. Note that these values are based on TSP samples collected by individual state agencies to comply with EPA Pb air monitoring regulations. The values of our data for site F, as seen in Figs. 3 and 4, are in general agreement with the Pb levels reported by the EPA. Our site F corresponds most closely to the urban core sites used by TCEQ for Pb monitoring.

3.2.2. Source of Lead

The order-of-magnitude higher concentrations of lead in the PM_{10} relative to those in the $\text{PM}_{2.5}$ is striking. Anthropogenic lead sources, e.g., the automobile emissions formerly associated with gasoline containing tetra-ethyl lead and related compounds or powdering old Pb-based paints, typically are associated with the fine fraction of airborne PM. Here the presence of the overwhelming majority of the lead in the PM_{10} fraction indicates the lead does not have an obvious current industrial, automotive, or other anthropogenic source. This observation is consistent with an earlier investigation of the speciation of lead in El Paso PM. Using synchrotron-based X-ray absorption spectroscopy (XAS), [20] demonstrated that virtually all the airborne lead was in the same chemical form as that in the local soils, i.e., adsorbed to humic acids or possibly such other soil components as clay particles. Thus the current airborne lead is a legacy of earlier pollution from automobile exhaust, a local lead, zinc, and copper smelter (shuttered in 1991), and other anthropogenic

sources. The speciation and the particle size of that lead changed with deposition and transformative chemical interactions within the local soil.

3.3. Geographic Trends

Sites F, H, and J stand out, in sequence, with the highest lead PM values. Site F is within 1.5 km of the former ASARCO smelter, adjacent to the University of Texas at El Paso, which is built on land donated by ASARCO in the early 20th Century. The smelter was founded in 1887 to handle lead ores from the mines in northern Mexico, and later added copper and zinc separations. It operated for more than a century, and ceased operations in 1999. The US EPA found ASARCO a potentially responsible party for lead and arsenic contamination in the neighborhoods surrounding the plant and undertook a cleanup of residential soils at some 1000 houses within a 5 km distance of the plant. In addition to soil contamination from the smelter, site F lies close to Interstate Highway 10, Mesa Street—State Highway 20 (a heavily trafficked commercial route), two railroads, and US Highway 62—Paisano Drive. Thus leaded gasoline and railroad transport also contributed to a significant legacy burden of soil contamination in this older part of El Paso's urban core [18,19,30]. Inasmuch as chemical speciation indicates that re-entrained soil is the dominant source of lead in current El Paso air [20], it is not surprising that the highest lead levels are coterminous with the part of the city bearing the most lead-contaminated soil.

Site H is in the Chamizal Memorial Monument, a small park located within 150 m of the US Customs and Border Patrol Inspection Station at the foot of the International Bridge of the Americas, a border crossing between El Paso, Texas, USA and Cd. Juarez, Chihuahua, Mexico. Some 400 m past the Customs Station is US Highway 54. On two other sides the small park borders US Highway 62 (50 m distant) and US Interstate Highway Loop 375—Cesar E. Chavez Border Highway (150 m). In addition to the legacy of airborne lead deposition onto the soil from vehicular traffic, the extremely busy bridge crossing has long been well known for traffic delays in excess of one or

two hours for entry into the United States. All vehicles are stopped at multiple lanes in a toll-booth-type barricade and drivers and passengers subjected to a brief—or extensive—questioning and vehicle scan or search. This process often causes backups of the multi-lane traffic, with motors running, extending a kilometer or more across the bridge over the Rio Grande into Mexico. Fallout from leaded gasoline was intense, and made more so by the Mexican vehicle fleet that was older and less well tuned than its US counterpart. Further, leaded gasoline was available in Mexico for some period of years after its regulation in the US. Not surprisingly, lead contamination of the soil in this area was high and thus we find high lead PM concentrations due to re-entrainment [19].

High airborne lead levels at Station J seem at first anomalous since this is a lower-population-density region. However, the sampling station lies a few hundred meters between two older thoroughfares, Socorro Road and Alameda Avenue—State Highway 20, the latter a main commercial route. The site is less than a kilometer from a large open-air trap and skeet range, a potential source of airborne lead fallout. In addition, westerly winds, unblocked by any topographic features, drive urban dust from Cd. Juarez into this area, yielding consistently high PM readings.

Site K bears similarities to site J. It is located proximal to, and downwind of, the Mexican border, adjacent to Border Highway—Interstate Highway. Lead levels are toward the middle range of those at our sites.

Lower levels of lead were recorded at the remaining stations. Sites A, D, I, and L are in residential areas of the city, with no special soils with high levels of legacy lead available for re-entrainment. Sites D and L, despite highways within a kilometer of each, exhibit particularly low levels of lead, consistent with their peripheral locations relative to urban development and its associated vehicular traffic during the bygone leaded-gasoline era.

3.4. Seasonal Trends

In general, the highest levels of airborne lead are associated with the winter season for PM_C and, to a lesser degree, for PM_F . Colder weather in El Paso often is accompanied by inversions that trap PM in a stagnant layer of air ranging from 200-500 m thickness above ground surface [31]. The higher PM loadings in winter result in higher lead values as well. Although strong winds occur throughout the year, spring is regarded as the windy season in El Paso, with occasional dust storms. It also is dry, typically with less than a few cm of rainfall. Inasmuch as re-entrained soil is locally the dominant source of airborne lead, it is reasonable that we observe higher lead levels in spring than in summer due to the higher loading of PM in the air. Rainfall in the summer monsoon helps clear PM from the air and decrease re-entrainment of Pb-contaminated soil.

3.5. Public Health Implications

In terms of public health, effective annual exposure of the typical El Paso resident falls well below regulatory levels of concern, as elaborated earlier in section 3.2.1. But a more careful parsing of the data indicates that

effective exposure actually is much lower because most or virtually all of the lead is contained in the PM_C . It is well established that PM_C can penetrate into the deep lung and settle in the bronchioles and alveoli. However, PM_F is even more effective at penetrating the alveolus in the gas exchange region of the lung and thus these are classified as *respirable* particles. PM_F also includes the sub-micron particles, and nanoparticles that are capable of entering the bloodstream. Exposure to the same chemical form or compound of lead in finer particles would be expected to result in greater transfer of lead to the exposed individual due to: 1) longer pulmonary retention time of the more deeply respired fine particles; 2) more intimate contact of those particles with tissue and cells of the lung; and 3) higher surface-area-to-volume ratio of the finer particles, resulting in easier chemical reaction and elemental exchange with tissue (release and transfer of Pb to tissue).

4. Conclusions

Levels of airborne lead in El Paso, Texas are well within the limits established by US EPA standards. The lead is concentrated in the PM_C fraction, consistent with the previously established contemporary source of re-entrained soil. Soil contamination is a legacy of leaded gas use, over a century of emissions from a lead smelter within 2 km of downtown, powdering lead building paint, and other traditional commercial and industrial sources. Seasonally, lead levels are highest in the fall—winter due to atmospheric trapping by diurnal temperature inversions. Levels are lowest in summer, and intermediate during spring. The geospatial distribution of airborne Pb across El Paso reveals highest levels in older areas where soil lead levels are highest. Overall, airborne lead should not represent a significant health consequence to most of El Paso's populations. Further, concentration of the lead in the PM_C fraction suggests a lower bioavailability of inhaled lead than would occur if the same amount of lead were concentrated in the PM_F fraction, which would penetrate more deeply into the pulmonary system.

Acknowledgements

This work was supported by grant number S11 ES013339 from the National Institute of Environmental Health Sciences (NIEHS) and the National Institutes of Health (NIH). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIEHS or NIH. We appreciate the efforts and support of all the faculty and students who participated in this NIEHS ARCH Program. We also thank the TCEQ and EPWU for their cooperation in providing access to their property and electrical power for the siting of our air samplers.

References

- [1] Settle, D.M., Patterson, C.C., "Lead in albacore: Guide to lead pollution in Americans," *Science*, 207 (4436). 1167-1176, 1980.
- [2] National Research Council, *Lead in the Human Environment*. Washington, D.C.: National Academy Press, 1980.

- [3] Needleman HL, editor. Human Lead Exposure. Boca Raton, FL: CRC Press; 1992.
- [4] Bryce-Smith, D., Waldron, H.A., Landrigan, P.J., Whitworth, R.H., Baloh, R.W., Lansdown, R.G., Clayton, B.E., Graham, P.J., Delves, H.T., "Blood-lead levels, behavior, and intelligence," *The Lancet*, 303(7867). 1166-1168, 1974.
- [5] Davidson, C.I., Rabinowitz, M., "Lead in the environment: From sources to human receptors," *Human Lead Exposure*, H.L. Needleman HL, ed., pp. 65-86, Boca Raton: CRC, 1992.
- [6] Annett, J., Dirkle, J., Makuc, C., Nesse, J., Bayse, D., et al., "Chronological trend in blood lead levels between 1976 and 1980," *New England Journal Medicine*, 308. 1373-1377, 1983.
- [7] Needleman, H.L., Leviton, A., Bellinger, D., "Lead-associated intellectual deficits," *New England Journal Medicine*, 306. 367-368, 2002.
- [8] ATSDR (Agency for Toxic Substances Disease Registry) "The Nature and Extent of Lead Poisoning in Children in the United States: A Report to Congress," Atlanta: US Department of Health and Human Services, Public Health Service, ATSDR, 1998.
- [9] Bellinger, D.C., Needleman, H.L., Leviton, A., Waternaux, C., Rabinowitz, M.B., Nichols, M.L., "Early sensory-motor development and prenatal exposure to lead," *Neurobehavioral Toxicology and Teratology*, 6. 387-402, 1984.
- [10] CDC (Centers for Disease Control. "Lead," <https://www.cdc.gov/nceh/lead/>. [Accessed on 6/24/2017].
- [11] Amaya, M.A., Jolly, K.W., Pingitore, N.E., "Blood lead in the 21st Century: The sub-microgram challenge," *Journal Blood Medicine*, 1. 71-78, 2010.
- [12] US Environmental Protection Agency 2010 Lead (Pb) Air Monitoring Requirements & 2013 Method for Determination of Lead (Pb) in Total Suspended Particulate Matter Fact Sheets US Environmental Protection Agency (2008b) Fact Sheet - Final Revisions to the National Ambient Air Quality Standards for Lead. US Environmental Protection Agency 2008b Fact Sheet - Final Revisions to the National Ambient Air Quality Standards for Lead. Available: <https://www.epa.gov/lead-air-pollution/2010-lead-pb-air-monitoring-requirements-2013-method-determination-lead-pb-total>. [Accessed on 5/31/2017].
- [13] US Environmental Protection Agency National Ambient Air Quality Standards (NAAQS) for Lead (Pb) Fact Sheets and Additional Information. Available: <https://www.epa.gov/lead-air-pollution/national-ambient-air-quality-standards-naaqs-lead-pb-fact-sheets-and-additional>. [Accessed on 5/31/2017].
- [14] Landrigan, P.J., Gehlbach, S.H., Rosenblum, B.F., Shoults, J.M., Candelaria, R.M., Barthel, W.F., Liddle, J.A., Smerk, A.L., Staehling, N.W., Sanders, J-D., "Epidemic lead absorption near an ore smelter – The role of particulate lead," *New England Journal of Medicine*, 292(3). 123-129, 1975.
- [15] Perales, M., *Smelertown: Making and Remembering a Southwest Border Community*, University of North Carolina Press, Chapel Hill, 2010.
- [16] Sullivan, M., *Tainted Earth: Smelters, Public Health, and the Environment*, Rutgers University Press, New Brunswick (NJ), 2014.
- [17] Camacho Araujo, I.B., *Airborne Lead in El Paso, Texas, 1977-1999*, M.S. Thesis, The University of Texas at El Paso, 2011.
- [18] Barnes, B.B., Pingitore, N.E., Mackay, W., "Levels of Toxic Metals in El Paso Soils proximal to Three Potential Industrial Point Sources," *The U.S. Mexican Border Environment: Integrated Approach to Defining Particulate Matter Issues in the Paso del Norte Region*, SCERP Monograph Series, no. 12, R.M. Currey, K.E. Kelley, H.L.C. Meuzelaar, and A.F. Serafim, eds., Chapter V, Section 4, 152-161, 2005.
- [19] Elkekli, A.R., *Lead and Associated Heavy Metal Distribution, El Paso, Texas*, Ph.D. Dissertation, The University of Texas at El Paso, 2013.
- [20] Pingitore Jr, N.E., Clague, J., Amaya, M.A., Maciejewska, B., Reynoso, J.J., "Urban Airborne Lead: X-ray Absorption Spectroscopy Establishes Soil as Dominant Source," *Public Library of Science One (PLoS One)*, 4(4). e5019, 2009. <http://www.plosone.org/article/info:doi/10.1371/journal.pone.0005019>.
- [21] https://en.wikipedia.org/wiki/El_Paso,_Texas. [Accessed on 6/01/2017].
- [22] <https://www.tceq.texas.gov/airquality/airmod/data/site>. [Accessed on 6/01/2017].
- [23] EPA Method IO-2.2, Sampling of Ambient Air for PM10 Using an Andersen Dichotomous Sampler, *In Compendium of Methods for the Determination of Inorganic Compounds in Ambient Air*, EPA/625/R-96/010a, Environmental Protection Development, June 1999.
- [24] Li, W-W., Orquiz, R., Pingitore Jr., N.E., Garcia, J.H., Espino, T.T., Gardea-Torresdey, J., Chow, J., Watson, J.W., "Analysis of temporal and spatial dichotomous PM air samples in the El Paso-Cd. Juarez air quality basin," *Journal of Air and Waste Management Association*, 51, 1511-1560, 2001.
- [25] EPA Method IO-3.3 Determination of Metals in Ambient Particulate Matter Using X-Ray Fluorescence (XRF) Spectroscopy, *In Compendium of Methods for the Determination of Inorganic Compounds in Ambient Air*, EPA/625/R-96/010a, Environmental Protection Development, June 1999.
- [26] <https://www-s.nist.gov/srmors/browseMaterials.cfm?subkey=5&tableid=54>. [Accessed on 6/01/2017].
- [27] <http://www.micromatter.com/xrf.php>. [Accessed on 6/01/2017].
- [28] South Coast Air Quality Management District (California, USA), <http://www.aqmd.gov/>. [Accessed on 7/17/2017].
- [29] US EPA, Air Data: Air Quality Data Collected at Outdoor Monitors Across the US, <https://www.epa.gov/outdoor-air-quality-data>. [Accessed on 7/17/2017].
- [30] Olvera, H.A., Garcia, M., Li, W-W., Yang, H., Amaya, M.A., Myers, O., Burchiel, S.W., Berwick, M., Pingitore, Jr., N.E., "Principal Component Analysis Optimization of a PM_{2.5} Land Use Regression Model with Small Monitoring Network," *Science of the Total Environment*, 425. 27-34, 2012.
- [31] Webb, W.L., *Mesometeorology at El Paso*, Texas Western Press, The University of Texas at El Paso, 1971, pp. 33-36.