

# Histone Deacetylase Inhibitors: A Hope for Cancer Patients

Ahmed M. Kabel<sup>1,2,\*</sup>, Fahad H. Baali<sup>1</sup>

<sup>1</sup>Department of Clinical Pharmacy, College of Pharmacy, Taif University, Taif, Saudi Arabia

<sup>2</sup>Department of Pharmacology, Faculty of Medicine, Tanta University, Tanta, Egypt

\*Corresponding author: drakabel@gmail.com

**Abstract** Cancer is a disease caused by genetic and genomic alterations such as amplifications, translocations, deletions or point mutations. Also, cancer development is caused by epigenetic changes due to modifications such as post-translational histone acetylations that can alter DNA and chromatin structures without alterations in the DNA sequence. Histones are the primary protein component of chromatin. It plays an important role in the interactions between the nucleosomes and within the nucleosome itself. Histone deacetylase enzyme is found in the nucleus of the different body cells and plays a vital role in remodeling of chromatin structure and controlling gene expression. Also, it was found to play an important role in the development of cancer. Histone deacetylase inhibitors which are used mainly in psychiatry and neurology may represent a new hope for cancer therapy. They act by various mechanisms which may help in prevention and treatment of cancer. Their combination with various lines of cancer therapy may represent a new hope for cancer patients.

**Keywords:** histone deacetylase, hope, cancer, patients

**Cite This Article:** Ahmed M. Kabel, and Fahad H. Baali, "Histone Deacetylase Inhibitors: A Hope for Cancer Patients." *Journal of Cancer Research and Treatment*, vol. 3, no. 3 (2015): 34-36. doi: 10.12691/jcrt-3-3-1.

## 1. Introduction

Histone deacetylase inhibitors (HDAC inhibitors, HDIs) are a group of compounds that interfere with the function of histone deacetylase enzyme. HDIs have been in common use in psychiatry and neurology as mood stabilizers and anti-epileptics [1]. Recent studies have investigated them as possible treatments for cancers, parasitic infections and inflammatory diseases [2].

Normally, histone acetyl transferases (HAT) acetylate the lysine residues in core histones of DNA leading to a less compact and more transcriptionally active chromatin [3]. On the contrary, histone deacetylases (HDAC) remove the acetyl groups from the lysine residues leading to formation of a condensed and transcriptionally silenced chromatin. Reversible modification of the terminal tails of core histones represents the major epigenetic mechanism for remodeling higher-order chromatin structure and controlling gene expression. HDAC inhibitors (HDI) block this action resulting in hyperacetylation of histones, thereby affecting gene expression [4].

Histone deacetylase inhibitors are cytostatic agents that inhibit the proliferation of tumor cells by inducing arrest of the cell cycle, differentiation and/or apoptosis. They exert their anti-tumor effects through induction of expression of tumor suppressor genes, through modulating that the acetylation/deacetylation of histones and/or non-histone proteins such as transcription factors [5]. Histone acetylation and deacetylation play important roles in the modulation of chromatin topology and the regulation of gene transcription. Inhibition of histone deacetylase leads to accumulation of

hyperacetylated nucleosome core histones in most regions of chromatin but affects the expression of only a small subset of genes, leading to transcriptional activation of some genes, but repression of another genes. Non-histone proteins such as transcription factors are also targets for acetylation with varying functional effects. Acetylation enhances the activity of some transcription factors such as the tumor suppressor p53 and the erythroid differentiation factor GATA-1 but may repress transcriptional activity of others including T cell factor and the co-activator ACTR. Recent studies have shown that the estrogen receptor alpha (ERalpha) can be hyperacetylated in response to histone deacetylase inhibition, suppressing ligand sensitivity and regulating transcriptional activation by histone deacetylase inhibitors [6]. Acetylation may play an important regulatory role in diverse nuclear receptor signaling functions. A number of structurally diverse histone deacetylase inhibitors have shown potent antitumor efficacy. Several compounds are currently in early phase clinical development as potential treatments for solid and hematological cancers both as monotherapy and in combination with cytotoxics and differentiation agents [7,8].

HDIs should not be considered to act only on HDACs but also act on nonhistone transcription factors and transcriptional co-regulators. HDIs can alter the degree of acetylation nonhistone effector molecules and, therefore, increase or repress the transcription of genes by this mechanism [9].

## 2. Classification of HDAC

Histone deacetylases can be classified into four groups; class I which includes HDAC1, -2, -3 and -8 are related to

yeast RPD3 gene; class II which includes HDAC4, -5, -6, -7, -9 and -10 are related to yeast Hda1 gene; class III which are related to the Sir2 gene and include SIRT1-7 and class IV which contains only HDAC11 and has features of both Class I and II [10].

### 3. Classification of HDI

HDIs usually act on Class I and Class II HDACs by binding to the zinc-containing catalytic domain of the HDACs [11]. These HDIs include hydroxamic acids (e.g. trichostatin A), cyclic tetrapeptides (e.g. trapoxin B), depsipeptides, benzamides, electrophilic ketones, and the aliphatic acid compounds (e.g. phenylbutyrate and valproic acid). Second-generation HDIs include the hydroxamic acids vorinostat, belinostat, LAQ824, and panobinostat and the benzamides (e.g. entinostat, mocetinostat) [12]. Also, nicotinamide, dihydrocoumarin, naphthopyranone and 2-hydroxynaphthaldehydes are derivatives of NAD which have the ability to inhibit Class III HDACs [13].

### 4. Clinical Uses of HDIs

HDIs have a long history of use in psychiatry and neurology as mood stabilizers and anti-epileptics (e.g. valproic acid). Recently, HDIs are being studied as a mitigator for neurodegenerative diseases such as Alzheimer's disease and Huntington's disease [14]. Enhancement of memory formation is increased in mice given the HDIs sodium butyrate [15]. It was found that some cognitive deficits were restored in transgenic mice that have a model of Alzheimer's disease by nicotinamide, a competitive HDI of class III sirtuins [16].

The HDAC inhibitors are under investigation for treatment of human immunodeficiency virus (HIV). It was reported that HDIs can flush HIV from the reservoirs. After that, a separate vaccination to eliminate HIV allows the immune system to neutralize the virus. Givinostat is also under investigations for treatment of polycythemia vera, thrombocythemia and myelofibrosis. Also, HDIs are being studied for protection of the cardiac muscles in acute myocardial infarction [17].

### 5. HDIs and Cancer

The results of several studies reported that HDIs may have potent anticancer effect [8,18,19]. The exact mechanisms of this effect are unclear but the epigenetic pathways were proposed [20]. HDIs can induce p21 expression, a regulator of p53's tumor suppressor activity. HDACs are involved in the pathway by which the retinoblastoma protein (pRb) suppresses cell proliferation [21]. The pRb protein is a part of a complex that attracts HDACs to the chromatin so that it will deacetylate histones [22]. HDAC1 negatively regulates the cardiovascular transcription factor Kruppel-like factor 5 through direct interaction [23]. Estrogen is well-established as a mitogenic factor implicated in the tumorigenesis and progression of breast cancer via its binding to the estrogen receptor alpha (ER $\alpha$ ). Recent data indicated that

chromatin inactivation mediated by HDAC and DNA methylation is a critical component of ER $\alpha$  silencing in human breast cancer cells. HDIs were licensed for treatment of cutaneous T cell lymphoma and multiple myeloma [24].

Recently, Zhu et al. [25] reported that trichostatin A (TSA) and sodium valproate (VPA), which are HDIs, significantly repressed the proliferation of chondrosarcoma cells in a concentration-dependent manner. TSA arrested the cell cycle in G2/M phase and VPA arrested the cell cycle in G1 phase. The tumor growth was markedly suppressed in mice treated with both TSA and VPA. Moreover, Pang et al. [26] found that vorinostat, an HDAC inhibitor, had a potent anticancer effect through affecting chromatin interactions at MYC/PVT1 locus. Pompo et al. [27] reported that novel HDIs induced growth arrest, apoptosis and differentiation in sarcoma cancer stem cells. When tested in human osteosarcoma, rhabdomyosarcoma, and Ewing's sarcoma stem cells, the new HDIs increased acetyl-H3 and acetyl-tubulin levels and inhibited cancer stem cells growth by apoptosis induction. At non-toxic doses, they promoted osteogenic differentiation.

### 6. Combination of HDIs with Other Anticancer Agents

As single agents, treatment with HDIs has demonstrated limited clinical benefit for patients with solid tumors, prompting the investigation of novel treatment combinations with other anti-cancer agents [28]. The acceptable toxicity profile associated with HDAC inhibitor treatment permits a broad integration into currently approved chemotherapy regimens [29]. One of these combinations is with DNA damage-inducing therapies. HDIs synergistically enhance the growth inhibition and apoptosis of DNA-damaging agents and irradiation. This occurs through a HDAC inhibitor-mediated increase in chromatin accessibility and downregulation of DNA repair [28]. Also, HDIs were combined to topoisomerase inhibitors. Vorinostat, which is one of HDIs, was found to enhance the effect of topotecan and SN-38 (the metabolite of irinotecan) in small-cell lung cancer and glioblastoma cells in vitro [30,31]. The cytotoxic effects of the topoisomerase I inhibitor karenitecin were enhanced by the combination with valproic acid [32]. Moreover, the combination between valproic acid and methotrexate had a beneficial effect on transplantable tumor model [8]. Raha et al. [33] reported that combined histone deacetylase inhibition and tamoxifen induced apoptosis in tamoxifen-resistant breast cancer models by reversing Bcl-2 overexpression.

HDIs combination with radiotherapy has been shown in various cancer cell lines including breast, prostate, lung, colon and cervical cancer. Preclinical studies suggested that the most effective treatment schedule requires pretreatment of cancer cells with the HDIs followed by ionizing radiation [34]. When administered before radiotherapy, treatment of colorectal xenograft models with vorinostat resulted in significantly reduced tumor volume compared with treatment with radiotherapy alone [35].

## References

- [1] Mwakwari S, Patil V, Guerrant W, Oyelere A (2010). Macrocyclic Histone Deacetylase Inhibitors. *Current Topics in Medicinal Chemistry*; 10(14): 1423-40.
- [2] Patil V, Guerrant W, Chen PC, Gryder B, Benicewicz DB, Khan SI, et al. (2010). Antimalarial and antileishmanial activities of histone deacetylase inhibitors with triazole-linked cap group. *Bioorganic & Medicinal Chemistry*; 18: 415-25.
- [3] Thiagalingam SAM, Cheng K-H, Lee HJ, Mineva N, Thiagalingam A, Ponte JF (2003). Histone Deacetylases: Unique Players in Shaping the Epigenetic Histone Code. *Annals of the New York Academy of Sciences*; 983: 84-100.
- [4] Dokmanovic M, Clarke C, Marks PA (2007). Histone Deacetylase Inhibitors: Overview and Perspectives. *Molecular Cancer Research*; 5 (10): 981-9.
- [5] Chueh AC, Tse JW, Tögel L, Mariadason JM (2015). Mechanisms of Histone Deacetylase Inhibitor-Regulated Gene Expression in Cancer Cells. *Antioxid Redox Signal*; 23(1):66-84.
- [6] Gryder BE, Rood MK, Johnson KA, Patil V, Raftery ED, Yao L-P, et al. (2013). Histone Deacetylase Inhibitors Equipped with Estrogen Receptor Modulation Activity. *J Med Chem*; 56 (14): 5782-96.
- [7] Vigushin D, Coombes R (2004). Targeted Histone Deacetylase Inhibition for Cancer Therapy. *Current Cancer Drug Targets*; 4 (2): 205-18.
- [8] Kabel AM (2014). Effect of Combination between Methotrexate and Histone Deacetylase Inhibitors on Transplantable Tumor Model. *Am J Med Studies*; 2(1): 12-18.
- [9] Yang X-J, Seto E (2007). HATs and HDACs: From structure, function and regulation to novel strategies for therapy and prevention. *Oncogene*; 26 (37): 5310-8.
- [10] Miller TA, Witter DJ, Belvedere S (2003). Histone deacetylase inhibitors. *J Med Chem*; 46 (24): 5097-116.
- [11] Drummond DC, Noble CO, Kirpotin DB, Guo Z, Scott GK, Benz CC (2005). Clinical Development of Histone Deacetylase Inhibitors As Anticancer Agents. *Ann Rev Pharmacol Toxicol*; 45: 495-528.
- [12] Beckers T, Burkhardt C, Wieland H, Gimmnich P, Ciossek T, Maier T, et al. (2007). Distinct pharmacological properties of second generation HDAC inhibitors with the benzamide or hydroxamate head group. *Int J Cancer*; 121 (5): 1138-48.
- [13] Porcu M, Chiarugi A (2005). The emerging therapeutic potential of sirtuin-interacting drugs: From cell death to lifespan extension. *Trends in Pharmacological Sciences*; 26 (2): 94-103.
- [14] Hahnen E, Hauke J, Tränkle C, Eyüpoglu IY, Wirth B, Blümcke I (2008). Histone deacetylase inhibitors: Possible implications for neurodegenerative disorders. *Expert Opinion on Investigational Drugs*; 17 (2): 169-84.
- [15] Guan J-S, Haggarty SJ, Giacometti E, Dannenberg J-H, Joseph N, Gao J, et al. (2009). HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature*; 459(7243): 55-60.
- [16] Green KN, Steffan JS, Martinez-Coria H, Sun X, Schreiber SS, Thompson LM, et al. (2008). Nicotinamide Restores Cognition in Alzheimer's Disease Transgenic Mice via a Mechanism Involving Sirtuin Inhibition and Selective Reduction of Thr231-Phosphotau. *Journal of Neuroscience*; 28 (45): 11500-10.
- [17] Granger A, Abdullah I, Huebner F, Stout A, Wang T, Huebner T, et al. (2008). Histone deacetylase inhibition reduces myocardial ischemia-reperfusion injury in mice. *The FASEB Journal*; 22 (10): 3549-60.
- [18] Abdel-Rahman MN, Kabel AM (2012). Comparative study between the effect of methotrexate and valproic acid on solid Ehrlich tumour. *J Egypt Natl Canc Inst*; 24: 161-7.
- [19] Fourcade S, Ruiz M, Guilera C, Hahnen E, Brichta L, Naudi A, et al. (2010). Valproic acid induces antioxidant effects in X-linked adrenoleukodystrophy. *Human Molecular Genetics*; 19 (10): 2005-14.
- [20] Mack GS (2010). To selectivity and beyond. *Nature Biotechnology*; 28 (12): 1259-66.
- [21] Richon VM, Sandhoff TW, Rifkind RA, Marks PA (2000). Histone deacetylase inhibitor selectively induces p21WAF1 expression and gene-associated histone acetylation. *Proc Natl Acad Sci*; 97 (18): 10014-9.
- [22] Kouzarides T, Miska A, McCance EA, Reid DJ, Bannister JL, Kouzarides AJ (1998). Retinoblastoma protein recruits histone deacetylase to repress transcription. *Nature*; 391 (6667): 597-601.
- [23] Matsumura T, Suzuki T, Aizawa K, Munemasa Y, Muto S, Horikoshi M, et al. (2004). The deacetylase HDAC1 negatively regulates the cardiovascular transcription factor Krüppel-like factor 5 through direct interaction. *J Biol Chem*; 280 (13): 12123-9.
- [24] Zhang Z, Yamashita H, Toyama T, Sugiura H, Ando Y, Mita K, et al. (2005). Quantitation of HDAC1 mRNA Expression in Invasive Carcinoma of the Breast. *Breast Cancer Research and Treatment*; 94 (1): 11-6.
- [25] Zhu J, Gu J, Ma J, Xu Z, Tao H (2015). Histone deacetylase inhibitors repress chondrosarcoma cell proliferation. *JBUON*; 20(1): 269-74.
- [26] Pang Q, Chee Y, Fullwood M (2015). Vorinostat, an HDAC inhibitor, perturbs chromatin interactions at MYC/PVT1 locus. *Front Genet*; 6:270.
- [27] Pompo GD, Salerno M, Rotili D, Valente S, Zwergel C, Avnet S, et al. (2015). Novel Histone Deacetylase Inhibitors Induce Growth Arrest, Apoptosis, and Differentiation in Sarcoma Cancer Stem Cells. *J Med Chem*.
- [28] Thurn KT, Thomas S, Moore A, Munster PN (2011). Rational therapeutic combinations with histone deacetylase inhibitors for the treatment of cancer. *Future Oncol*; 7(2): 263-83.
- [29] Munster PN, Marchion D, Thomas S, et al. (2009). Phase I trial of vorinostat and doxorubicin in solid tumours: histone deacetylase 2 expression as a predictive marker. *Br J Cancer*; 101(7): 1044-50.
- [30] Bruzzese F, Rocco M, Castelli S, Di Gennaro E, Desideri A, Budillon A (2009). Synergistic antitumor effect between vorinostat and topotecan in small cell lung cancer cells is mediated by generation of reactive oxygen species and DNA damage-induced apoptosis. *Mol Cancer Ther*; 8(11): 3075-87.
- [31] Sarcar B, Kahali S, Chinnaiyan P (2010). Vorinostat enhances the cytotoxic effects of the topoisomerase I inhibitor Sn38 in glioblastoma cell lines. *J Neurooncol*; 99(2): 201-7.
- [32] Daud AI, Dawson J, Deconti RC, et al. (2009). Potentiation of a topoisomerase I inhibitor, karenitecin, by the histone deacetylase inhibitor valproic acid in melanoma: translational and Phase I/II clinical trial. *Clin Cancer Res*; 15(7): 2479-87.
- [33] Raha P, Thomas S, Thurn KT, Park J, Munster PN (2015). Combined histone deacetylase inhibition and tamoxifen induces apoptosis in tamoxifen-resistant breast cancer models, by reversing Bcl-2 overexpression. *Breast Cancer Research*; 17: 26.
- [34] Nome RV, Bratland A, Harman G, Fodstad O, Andersson Y, Ree AH (2005). Cell cycle checkpoint signaling involved in histone deacetylase inhibition and radiation-induced cell death. *Mol Cancer Ther*; 4(8): 1231-8.
- [35] Folkvord S, Ree AH, Furre T, Halvorsen T, Flatmark K (2009). Radiosensitization by SAHA in experimental colorectal carcinoma models – in vivo effects and relevance of histone acetylation status. *Int J Radiat Oncol Biol Phys*; 74(2): 546-52.