

# q-Hardy-Littlewood-Type Maximal Operator with Weight Related to Fermionic p-Adic q-Integral on $Z_p$

Erdoğan Şen<sup>1</sup>, Mehmet Acikgoz<sup>2</sup>, Serkan Araci<sup>3,\*</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science and Letters, Namik Kemal University, Tekirdağ, Turkey

<sup>2</sup>Department of Mathematics, Faculty of Science and Arts, University of Gaziantep, Gaziantep, Turkey

<sup>3</sup>Atatürk Street, 31290 Hatay, TURKEY

\*Corresponding author: mtsrkn@hotmail.com

Received August 14, 2013; Revised September 17, 2013; Accepted September 25, 2013

**Abstract** The q-extension of Hardy-littlewood-type maximal operator in accordance with q-Volkenborn integral in the p-adic integer ring was recently studied [11]. A generalization of Jang's results was given by Araci and Acikgoz [1]. By the same motivation of their papers, we aim to give the definition of the weighted q-Hardy-littlewood-type maximal operator by means of fermionic p-adic q-invariant distribution on  $Z_p$ . Finally, we derive some interesting properties involving this-type maximal operator.

**Keywords:** fermionic p-adic q-integral on  $Z_p$ , hardy-littlewood theorem, p-adic analysis, q-analysis

**Cite This Article:** Erdoğan Şen, Mehmet Acikgoz, and Serkan Araci, "q-Hardy-Littlewood-Type Maximal Operator with Weight Related to Fermionic p-Adic q-Integral on  $Z_p$ ." *Turkish Journal of Analysis and Number Theory* 1, no. 1 (2013): 4-8. doi: 10.12691/tjant-1-1-2.

## 1. Introduction

The concept of p-adic numbers was originally invented by Kurt Hensel who is German mathematician, around the end of the nineteenth century [12]. In spite of their being already one hundred years old, these numbers are still today enveloped in an aura of mystery within the scientific community and also play a vital and important role in mathematics.

The fermionic p-adic q-integral in the p-adic integer ring was originally constructed by Kim [2,6] who introduced Lebesgue-Radon-Nikodym Theorem with respect to fermionic p-adic q-integral on  $Z_p$ . The fermionic p-adic q-integral on  $Z_p$  is used in mathematical physics for example the functional equation of the q-zeta function, the q-stirling numbers and q-mahler theory of integration with respect to the ring  $Z_p$  together with Iwasawa's p-adic q-L function.

In [11], Jang defined q-extension of Hardy-Littlewood-type maximal operator by means of q-Volkenborn integral on  $Z_p$ . Afterwards, in [1], Araci and Acikgoz added a weight into Jang's q-Hardy-Littlewood-type maximal operator and derived some interesting properties by means of Kim's p-adic q-integral on  $Z_p$ . Now also, we shall consider weighted q-Hardy-Littlewood-type maximal operator on the fermionic p-adic q-integral on  $Z_p$ . Moreover, we shall analyse q-Hardy-Littlewood-type maximal operator via the fermionic p-adic q-integral on  $Z_p$ .

Assume that p be an odd prime number. Let  $Q_p$  be the field of p-adic rational numbers and let  $C_p$  be the completion of algebraic closure of  $Q_p$ .

Thus,

$$Q_p = \left\{ x = \sum_{n=-k}^{\infty} a_n p^n : 0 \leq a_n < p \right\}.$$

Then  $Z_p$  is an integral domain to be

$$Z_p = \left\{ x = \sum_{n=0}^{\infty} a_n p^n : 0 \leq a_n \leq p-1 \right\},$$

or

$$Z_p = \left\{ x \in Q_p : |x|_p \leq 1 \right\}.$$

In this paper, we assume that  $q \in C_p$  with  $|1-q|_p < 1$  as an indeterminate.

The p-adic absolute value  $|\cdot|_p$ , is normally defined by

$$|x|_p = \frac{1}{p^r},$$

where  $x = p^r \frac{s}{t}$  with  $(p,s) = (p,t) = (s,t) = 1$  and  $r \in \mathbb{Q}$ .

A p-adic Banach space  $B$  is a  $Q_p$ -vector space with a lattice  $B^0$  ( $Z_p$ -module) separated and complete for p-adic topology, ie.,

$$B^0 \simeq \varprojlim_{n \in \mathbb{N}} B^0 / p^n B^0.$$

For all  $x \in B$ , there exists  $n \in \mathbb{Z}$ , such that  $x \in p^n B^0$ .

Define

$$v_B(x) = \sup_{n \in \mathbb{N} \cup \{+\infty\}} \{n : x \in p^n B^0\}.$$

It satisfies the following properties:

$$v_B(x+y) \geq \min(v_B(x), v_B(y)),$$

$$v_B(\beta x) = v_p(\beta) + v_B(x), \text{ if } \beta \in \mathbb{Q}_p.$$

Then,  $\|x\|_B = p^{-v_B(x)}$  defines a norm on  $B$ , such that  $B$  is complete for  $\|\cdot\|_B$  and  $B^0$  is the unit ball.

A measure on  $Z_p$  with values in a p-adic Banach space  $B$  is a continuous linear map

$$f \mapsto \int f(x) \mu = \int_{Z_p} f(x) \mu(x)$$

from  $C^0(Z_p, C_p)$ , (continuous function on  $Z_p$ ) to  $B$ . We know that the set of locally constant functions from  $Z_p$  to  $\mathbb{Q}_p$  is dense in  $C^0(Z_p, C_p)$  so.

Explicitly, for all  $f \in C^0(Z_p, C_p)$ , the locally constant functions

$$f_n = \sum_{i=0}^{p^n-1} f(i) 1_{i+p^n Z_p} \rightarrow f \text{ in } C^0.$$

Now if  $\mu \in \mathfrak{D}_0(Z_p, \mathbb{Q}_p)$ , set  $\mu(i+p^n Z_p) = \int_{Z_p} 1_{i+p^n Z_p} \mu$ .

Then  $\int_{Z_p} f \mu$  is given by the following Riemann sums

$$\int_{Z_p} f \mu = \lim_{n \rightarrow \infty} \sum_{i=0}^{p^n-1} f(i) \mu(i+p^n Z_p).$$

T. Kim defined  $\mu_{-q}$  as follows:

$$\mu_{-q}(\xi + dp^n Z_p) = \frac{(-q)^\xi}{[dp^n]_{-q}}$$

and this can be extended to a distribution on  $Z_p$ . This distribution yields an integral in the case  $d = 1$ .

So, q-Volkenborn integral was defined by T. Kim as follows:

$$I_{-q}(f) = \int_{Z_p} f(\xi) d\mu_q(\xi)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{[p^n]_{-q}} \sum_{\xi=0}^{p^n-1} (-1)^\xi f(\xi) q^\xi \tag{1.1}$$

where  $[x]_q$  is a q-extension of  $x$  which is defined by

$$[x]_q = \frac{1-q^x}{1-q}.$$

Note that  $\lim_{q \rightarrow 1} [x]_q = x$  cf. [1,2,4,5,6,7,11].

Let  $d$  be a fixed positive integer with  $(p, d) = 1$ . We now set

$$X = X_d = \varinjlim_n Z / dp^n Z,$$

$$X_1 = Z_p,$$

$$X^* = \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} a + dpZ_p,$$

$$a + dp^n Z_p = \{x \in X \mid x \equiv a \pmod{p^n}\},$$

where  $a \in \mathbb{Z}$  satisfies the condition  $0 \leq a < dp^n$ . For  $f \in UD(Z_p, C_p)$ ,

$$\int_{Z_p} f(x) d\mu_{-q}(x) = \int_X f(x) d\mu_{-q}(x) \text{ see [10]}$$

By means of q-Volkenborn integral, we consider below strongly p-adic q-invariant distribution  $\mu_{-q}$  on  $Z_p$  in the form

$$\left| \begin{matrix} [p^n]_{-q} \mu_{-q}(a + p^n Z_p) \\ -[p^{n+1}]_{-q} \mu_{-q}(a + p^{n+1} Z_p) \end{matrix} \right| < \delta_n,$$

where  $\delta_n \rightarrow 0$  as  $n \rightarrow \infty$  and  $\delta_n$  is independent of  $a$ . Let  $f \in UD(Z_p, C_p)$ , for any  $a \in Z_p$ , we assume that the weight function  $\omega(x)$  is defined by  $\omega(x) = \omega^x$  where  $\omega \in C_p$  with  $|1-\omega|_p < 1$ . We define the weighted measure on  $Z_p$  as follows:

$$\mu_{f,-q}^{(\omega)}(a + p^n Z_p) = \int_{a+p^n Z_p} \omega^\xi f(\xi) d\mu_{-q}(\xi) \tag{1.2}$$

where the integral is the fermionic p-adic q-integral on  $Z_p$ . From (1.2), we note that  $\mu_{f,-q}^{(\omega)}$  is a strongly weighted measure on  $Z_p$ . Namely,

$$\left| \begin{matrix} [p^n]_{-q} \mu_{f,-q}^{(\omega)}(a + p^n Z_p) \\ -[p^{n+1}]_{-q} \mu_{f,-q}^{(\omega)}(a + p^{n+1} Z_p) \end{matrix} \right|_p$$

$$= \left| \sum_{x=0}^{p^n-1} (-1)^x \omega^x f(x) q^x - \sum_{x=0}^{p^{n+1}-1} (-1)^x \omega^x f(x) q^x \right|_p$$

$$\leq \left| \frac{f(p^n) (-1)^{p^n} \omega^{p^n} q^{p^n}}{p^n} \right|_p \left| p^n \right|_p$$

$$\leq Cp^{-n}$$

Thus, we get the following proposition.

**Proposition 1.** For  $f, g \in UD(Z_p, C_p)$ , then, we have

$$\begin{aligned} & \mu_{\alpha f + \beta g, -q}^{(\omega)}(a + p^n Z_p) \\ &= \alpha \mu_{f, -q}^{(\omega)}(a + p^n Z_p) + \beta \mu_{g, -q}^{(\omega)}(a + p^n Z_p). \end{aligned}$$

where  $\alpha, \beta$  are positive constants. Also, we have

$$\left| \begin{array}{l} [p^n]_{-q} \mu_{f, -q}^{(\omega)}(a + p^n Z_p) \\ - [p^{n+1}]_{-q} \mu_{f, -q}^{(\omega)}(a + p^{n+1} Z_p) \end{array} \right| \leq Cp^{-n}$$

where  $C$  is positive constant.

Let  $\mathbf{P}_q(x) \in C_p[[x]_q]$  be an arbitrary  $q$ -polynomial.

Now also, we indicate that  $\mu_{\mathbf{P}, -q}^{(\omega)}$  is a strongly weighted fermionic  $p$ -adic  $q$ -invariant measure on  $Z_p$ . Without a loss of generality, it is sufficient to evidence the statement for  $\mathbf{P}(x) = [x]_q^k$ .

$$\begin{aligned} & \mu_{\mathbf{P}, -q}^{(\omega)}(a + p^n Z_p) \\ &= \lim_{m \rightarrow \infty} \frac{1}{[p^m]_{-q}} \sum_{i=0}^{p^{m-n}-1} w^{a+ip^n} [a+ip^n]_q^k (-q)^{a+ip^n} \end{aligned} \quad (1.3)$$

where

$$\begin{aligned} & [a+ip^n]_q^k \\ &= \sum_{j=0}^k \binom{k}{j} [a]_q^{k-j} q^{aj} [p^n]_q^j [i]_q^j p^n \\ &= [a]_q^k + k [a]_q^{k-1} q^a [p^n]_q [i]_q p^n \\ & \quad + \dots + q^{ak} [p^n]_q^k [i]_q^k p^n \end{aligned} \quad (1.4)$$

and

$$\begin{aligned} w^{a+ip^n} &= w^a \sum_{l=0}^{ip^n} \binom{ip^n}{l} (w-1)^l \\ &\equiv w^a \pmod{p^n}. \end{aligned} \quad (1.5)$$

By (1.5), we have

$$\begin{aligned} & (-q)^{a+ip^n} \\ &= (-q)^a \sum_{l=0}^{ip^n} \binom{ip^n}{l} (-1)^l (q+1)^l \\ &\equiv (-q)^a \pmod{p^n}. \end{aligned} \quad (1.6)$$

By (1.3), (1.4), (1.5) and (1.6), we have the following

$$\begin{aligned} & \mu_{\mathbf{P}, -q}^{(\omega)}(a + p^n Z_p) \\ &\equiv (-1)^a \omega^a q^a [a]_q^k \pmod{p^n} \\ &\equiv (-1)^a \omega^a q^a \mathbf{P}(a) \pmod{p^n}. \end{aligned}$$

For  $x \in Z_p$ , let  $x \equiv x_n \pmod{p^n}$  and  $x \equiv x_{n+1} \pmod{p^{n+1}}$ , where  $x_n, x_{n+1} \in Z$  with  $0 \leq x_n < p^n$  and  $0 \leq x_{n+1} < p^{n+1}$

Then, we procure the following

$$\left| \begin{array}{l} [p^n]_{-q} \mu_{\mathbf{P}, -q}^{(\omega)}(a + p^n Z_p) \\ - [p^{n+1}]_{-q} \mu_{\mathbf{P}, -q}^{(\omega)}(a + p^{n+1} Z_p) \end{array} \right| \leq Cp^{-n},$$

where  $C$  is positive constant and  $n \gg 0$ .

Let  $UD(Z_p, C_p)$  be the space of uniformly differentiable functions on  $Z_p$  with sup-norm

$$\|f\|_\infty = \sup_{x \in Z_p} |f(x)|_p.$$

The difference quotient  $\Delta_1 f$  of  $f$  is the function of two variables given by

$$\Delta_1 f(m, x) = \frac{f(x+m) - f(x)}{m}$$

for all  $x, m \in Z_p, m \neq 0$

A function  $f : Z_p \rightarrow C_p$  is said to be a Lipschitz function if there exists a constant  $M > 0$  (the Lipschitz constant of  $f$ ) such that

$$|\Delta_1 f(m, x)| \leq M \text{ for all } m \in Z_p \setminus \{0\} \text{ and } x \in Z_p.$$

The  $C_p$  linear space consisting of all Lipschitz function is denoted by  $Lip(Z_p, C_p)$ . This space is a Banach space with the respect to the norm  $\|f\|_1 = \|f\|_\infty \vee \|\Delta_1 f\|_\infty$  (for more information, see [3-9]). The objective of this paper is to introduce weighted  $q$ -Hardy Littlewood-type maximal operator on the fermionic  $p$ -adic  $q$ -integral on  $Z_p$ . Also, we show that the boundedness of the weighted  $q$ -Hardy-littlewood-type maximal operator in the  $p$ -adic integer ring.

## 2. The Weighted $q$ -Hardy-Littlewood-Type Maximal Operator

In view of (1.2) and the definition of fermionic  $p$ -adic  $q$ -integral on  $Z_p$ , we now consider the following theorem.

**Theorem 1.** Let  $\mu_{-q}^{(w)}$  be a strongly fermionic  $p$ -adic  $q$ -invariant on  $Z_p$  and  $f \in UD(Z_p, C_p)$ . Then for any  $n \in Z$  and any  $\xi \in Z_p$ , we have

$$\begin{aligned} & \int_{a+p^n Z_p} \omega^\xi f(\xi) (-q)^{-\xi} d\mu_{-q}(\xi) \\ (1) &= \frac{(-1)^a \omega^a}{[p^n]_{-q}} \int_{Z_p} \omega^\xi f(a + p^n \xi) (-q)^{-p^n \xi} d\mu_{-q}(\xi) \end{aligned}$$

$$(2) \int_{a+p^n Z_p} \omega^\xi d\mu_{-q}(\xi) = \frac{\omega^a (-q)^a}{\left[ p^n \right]_{-q}} \frac{2}{1 + \omega^{p^n} q^{p^n}}$$

**Proof.** (1) By using (1.1) and (1.2), we see the following applications:

$$\begin{aligned} & \int_{a+p^n Z_p} \omega^\xi f(\xi) (-q)^{-\xi} d\mu_{-q}(\xi) \\ &= \lim_{m \rightarrow \infty} \frac{1}{\left[ p^{m+n} \right]_{-q}} \sum_{\xi=0}^{p^m-1} \left[ \begin{array}{l} \omega^{a+p^n \xi} f(a+p^n \xi) \\ \times (-q)^{-(a+p^n \xi)} \\ \times q^{a+p^n \xi} (-1)^{a+p^n \xi} \end{array} \right] \\ &= (-1)^a \omega^a \lim_{m \rightarrow \infty} \left[ \begin{array}{l} \frac{1}{\left[ p^m \right]_{-q} p^n \left[ p^n \right]_{-q}} \\ \times \sum_{\xi=0}^{p^m-1} \omega^\xi (-q)^{-p^n \xi} \\ \times f(a+p^n \xi) \left( -q^{p^n} \right)^\xi \end{array} \right] \\ &= \frac{(-1)^a \omega^a}{\left[ p^n \right]_{-q}} \int_{Z_p} \left[ \begin{array}{l} \omega^\xi f(a+p^n \xi) \\ \times (-q)^{-p^n \xi} d\mu_{-q^{p^n}}(\xi) \end{array} \right]. \end{aligned}$$

(2) By the same method of (1), then, we easily derive the following

$$\begin{aligned} & \int_{a+p^n Z_p} \omega^\xi d\mu_{-q}(\xi) \\ &= \lim_{m \rightarrow \infty} \frac{1}{\left[ p^{m+n} \right]_{-q}} \sum_{\xi=0}^{p^m-1} \omega^{a+\xi p^n} (-q)^{a+\xi p^n} \\ &= \frac{\omega^a (-q)^a}{\left[ p^n \right]_{-q}} \lim_{m \rightarrow \infty} \frac{1}{\left[ p^m \right]_{-q} p^n} \sum_{\xi=0}^{p^m-1} \left( \omega^{p^n} \right)^\xi \left( -q^{p^n} \right)^\xi \\ &= \frac{\omega^a (-q)^a}{\left[ p^n \right]_{-q}} \lim_{m \rightarrow \infty} \frac{1 + \left( \omega^{p^n} q^{p^n} \right)^{p^m}}{1 + \omega^{p^n} q^{p^n}} \\ &= \frac{\omega^a (-q)^a}{\left[ p^n \right]_{-q}} \frac{2}{1 + \omega^{p^n} q^{p^n}} \end{aligned}$$

Since  $\lim_{m \rightarrow \infty} q^{p^m} = 1$  for  $|1-q|_p < 1$ , our assertion follows.

We are now ready to introduce the definition of the weighted q-Hardy-littlewood-type maximal operator related to fermionic p-adic q-integral on  $Z_p$  with a strong fermionic p-adic q-invariant distribution  $\mu_{-q}$  in the p-adic integer ring.

**Definition 1.** Let  $\mu_{-q}^{(\omega)}$  be a strongly fermionic p-adic q-invariant distribution on  $Z_p$  and  $f \in UD(Z_p, C_p)$ . Then, q-Hardy-littlewood-type maximal operator with weight related to fermionic p-adic q-integral on  $a+p^n Z_p$  is defined as

$$\begin{aligned} & \mathbf{M}_{p,q}^{(\omega)} f(a) \\ &= \sup_{n \in Z} \frac{1}{\mu_{1,-q}^{(\omega)}(\xi + p^n Z_p)} \int_{a+p^n Z_p} \omega^\xi (-q)^{-\xi} f(\xi) d\mu_{-q}(\xi) \end{aligned}$$

for all  $a \in Z_p$ .

We recall that famous Hardy-littlewood maximal operator  $\mathbf{M}_\mu$ , which is defined by

$$\mathbf{M}_\mu f(a) = \sup_{a \in Q} \frac{1}{\mu(Q)} \int_Q |f(x)| d\mu(x), \quad (2.1)$$

where  $f : \mathbb{R}^k \rightarrow \mathbb{R}^k$  is a locally bounded Lebesgue measurable function,  $\mu$  is a Lebesgue measure on  $(-\infty, \infty)$  and the supremum is taken over all cubes  $Q$  which are parallel to the coordinate axes. Note that the boundedness of the Hardy-Littlewood maximal operator serves as one of the most important tools used in the investigation of the properties of variable exponent spaces (see [11]). The essential aim of Theorem 1 is to deal mainly with the weighted q-extension of the classical Hardy-Littlewood maximal operator in the space of p-adic Lipschitz functions on  $Z_p$  and to find the boundedness of them. By means of Definition 1, then, we state the following theorem.

**Theorem 2.** Let  $f \in UD(Z_p, C_p)$  and  $x \in Z_p$ , we get

$$(1) \quad \mathbf{M}_{p,q}^{(\omega)} f(a) = \frac{(-1)^a}{2q^a} \sup_{n \in Z} \left( 1 + \omega^{p^n} q^{p^n} \right) \int_{Z_p} \omega^\xi f(x + p^n \xi) (-q)^{-p^n \xi} d\mu_{-q^{p^n}}(\xi)$$

$$(2) \quad \left| \mathbf{M}_{p,q}^{(\omega)} f(a) \right|_p \leq \left| \frac{(-1)^a}{2q^a} \right|_p \sup_{n \in Z} \left| 1 + \omega^{p^n} q^{p^n} \right|_p \|f\|_1 \left\| \left( \frac{-q^{p^n}}{\omega} \right)^{-(\cdot)} \right\|_{L^1}$$

$$\text{where } \left\| \left( \frac{-q^{p^n}}{\omega} \right)^{-(\cdot)} \right\|_{L^1} = \int_{Z_p} \left( \frac{-q^{p^n}}{\omega} \right)^{-\xi} d\mu_{-q^{p^n}}(\xi).$$

**Proof.** (1) Because of Theorem 1 and Definition 1, we see

$$\begin{aligned} \mathbf{M}_{p,q}^{(\omega)} f(a) &= \sup_{n \in Z} \frac{1}{\mu_{1,-q}^{(\omega)}(\xi + p^n Z_p)} \\ & \int_{a+p^n Z_p} \omega^\xi (-q)^{-\xi} f(\xi) d\mu_{-q}(\xi) \end{aligned}$$

$$= \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left( 1 + \omega^{p^n} q^{p^n} \right) \int_{Z_p} \omega^\xi f(x + p^n \xi) (-q)^{-p^n \xi} d\mu_{-q^{p^n}}(\xi).$$

(2) On account of (1), we can derive the following

$$\begin{aligned} & \left| \mathbf{M}_{p,q}^{(\omega)} f(a) \right|_p \\ &= \left| \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left( 1 + \omega^{p^n} q^{p^n} \right) \int_{Z_p} \omega^\xi f(x + p^n \xi) (-q)^{-p^n \xi} d\mu_{-q^{p^n}}(\xi) \right|_p \\ &\leq \left| \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left( 1 + \omega^{p^n} q^{p^n} \right) \int_{Z_p} \omega^\xi f(x + p^n \xi) (-q)^{-p^n \xi} d\mu_{-q^{p^n}}(\xi) \right|_p \\ &\leq \left| \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left| 1 + \omega^{p^n} q^{p^n} \right|_p \int_{Z_p} |f(a + p^n \xi)|_p \left| \left( \frac{-q^{p^n}}{\omega} \right)^{-\xi} \right|_p d\mu_{-q^{p^n}}(\xi) \right|_p \\ &\leq \left| \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left| 1 + \omega^{p^n} q^{p^n} \right|_p \|f\|_1 \int_{Z_p} \left| \left( \frac{-q^{p^n}}{\omega} \right)^{-\xi} \right|_p d\mu_{-q^{p^n}}(\xi) \right|_p \\ &= \left| \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left| 1 + \omega^{p^n} q^{p^n} \right|_p \|f\|_1 \left\| \left( \frac{-q^{p^n}}{\omega} \right)^{-\cdot} \right\|_{L^1} \right|_p. \end{aligned}$$

Thus, we complete the proof of theorem.

We note that Theorem 2 (2) shows the supnorm-inequality for the q-Hardy-Littlewood-type maximal operator with weight on  $Z_p$ , on the other hand, Theorem 2 (2) shows the following inequality

$$\begin{aligned} & \left\| \mathbf{M}_{p,q}^{(\omega)} f \right\|_\infty \\ &= \sup_{x \in Z_p} \left| \mathbf{M}_{p,q}^{(\omega)} f(x) \right|_p \tag{2.2} \\ &\leq \mathbf{K} \|f\|_1 \left\| \left( \frac{-q^{p^n}}{\omega} \right)^{-\cdot} \right\|_{L^1} \end{aligned}$$

where  $\mathbf{K} = \left| \frac{(-1)^a}{2q^a} \sup_{n \in \mathbb{Z}} \left| 1 + \omega^{p^n} q^{p^n} \right|_p \right|_p$ . By the equation

(2.2), we get the following Corollary, which is the boundedness for weighted q-Hardy-Littlewood-type maximal operator with weight on  $Z_p$ .

**Corollary 1.**  $\mathbf{M}_{p,q}^{(\omega)}$  is a bounded operator from  $UD(Z_p, C_p)$  into  $L^\infty(Z_p, C_p)$ , where  $L^\infty(Z_p, C_p)$  is the space of all p-adic supnorm-bounded functions with the

$$\|f\|_\infty = \sup_{x \in Z_p} |f(x)|_p,$$

for all  $f \in L^\infty(Z_p, C_p)$ .

### References

- [1] S. Araci and M. Acikgoz, A note on the weighted q-Hardy-littlewood-type maximal operator with respect to q-Volkenborn integral in the p-adic integer ring, *J. Appl. Math. & Informatics*, Vol. 31(2013), No. 3-4, pp. 365-372.
- [2] S. Araci, M. Acikgoz and E. Şen, On the extended Kim's p-adic q-deformed fermionic integrals in the p-adic integer ring, *Journal of Number Theory* 133 (2013) 3348-3361.
- [3] T. Kim, Lebesgue-Radon-Nikodym theorem with respect to fermionic p-adic invariant measure on  $Z_p$ , *Russ. J. Math. Phys.* 19 (2012).
- [4] T. Kim, Lebesgue-Radon-Nikodym theorem with respect to fermionic q-Volkenborn distribution on  $\mu_p$ , *Appl. Math. Comp.* 187 (2007), 266-271.
- [5] T. Kim, S. D. Kim, D.W. Park, On Uniformly differntiability and q-Mahler expansion, *Adv. Stud. Contemp. Math.* 4 (2001), 35-41.
- [6] T. Kim, q-Volkenborn integration, *Russian J. Math. Phys.* 9 (2002) 288-299.
- [7] T. Kim, On a q-analogue of the p-adic log Gamma functions and related integrals, *Journal of Number Theory* 76 (1999), 320-329.
- [8] T. Kim, Note on Dedekind-type DC sums, *Advanced Studies in Contemporary Mathematics* 18(2) (2009), 249-260.
- [9] T. Kim, A note on the weighted Lebesgue-Radon-Nikodym Theorem with respect to p-adic invariant integral on  $Z_p$ , *J. Appl. Math. & Informatics*, Vol. 30(2012), No. 1, 211-217.
- [10] T. Kim, Non-archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials, *Russ. J. Math Phys.* 10 (2003) 91-98.
- [11] L-C. Jang, On the q-extension of the Hardy-littlewood-type maximal operator related to q -Volkenborn integral in the p-adic integer ring, *Journal of Chungcheon Mathematical Society*, Vol. 23, No. 2, June 2010.
- [12] K. Hensel, *Theorie der Algebraischen Zahlen I.* Teubner, Leipzig, 1908.
- [13] N. Koblitz, *p-adic Numbers, p-adic Analysis and Zeta Functions*, Springer-Verlag, New York Inc, 1977.